Quasigroups and Related Systems **33** (2025), 231 – 254

https://doi.org/10.56415/qrs.v33.18

Quasi-antiassociative bialgebras

Cyrille Essossolim Haliya, Gbêvèwou Damien Houndedji,

Bakary Kourouma

Abstract. This work explores key characteristics and properties of quasi-antiassociative algebras, including their bimodules, matched pairs and bialgebraic structures.

1. Introduction

Mock-Lie algebras, recently introduced by P. Zusmanovich in [15], are commutative algebras satisfying the Jacobi identity. These algebras have appeared in the literature under various names (see [15] and [4] for details). Two notable features of Mock-Lie algebras are: first, that algebras over the operad Koszul dual to the Mock-Lie operad can be characterized in three equivalent ways (see [15], [4]), and second, as noted in [13], Mock-Lie algebras can be derived from antiassociative algebras similarly to how they arise from associative algebras, indicating a strong link between antiassociative and Mock-Lie algebras. Here, "antiassociative algebras" refer to those where the operation $(a, b) \to ab$ satisfies $(ab)c+a(bc) = 0$ for each a, b, c. This class of algebras first appeared notably in [13], with main properties given, and in [10] where M. Markl and E. Remm (2014) established results on Koszul operads for n-ary algebras, with a focus on antiassociative operations.

In 2007, C. Bai examined left-symmetric bialgebras, which serve as analogues to Lie bialgebras, in [1]. Left-symmetric algebras (also called pre-Lie, quasi-associative, and Vinberg algebras) are Lie-admissible algebras where left multiplication operators form a Lie algebra; these were first introduced by Cayley in 1896. Bai highlighted that left-symmetric algebras also emerged in various studies in geometry and algebra, such as convex homogeneous cones [14, 9], affine manifolds, affine structures on Lie groups [11], and associative algebra deformation [6]. Notably, a Lie algebra $\mathcal{G}$ with

2010 Mathematics Subject Classification: 16T25, 05C25, 16S99, 16Z05.

Keywords: Quasi-antiassociative algebra, Mock-Lie (admissible) algebra, bialgebra

compatible left-symmetric structure is the Lie algebra of a Lie group $\mathcal{G}$ with a left-invariant affine structure, where there is a left-invariant, flat, and torsion-free connection $\bigtriangledown$ in G. The left-symmetric algebra structure corresponds to the connection $\bigtriangledown$ given by $XY = \bigtriangledown_X Y$ for $X, Y \in G$, where left-symmetry reflects the connection's flatness [8, 12].

Chapoton and Livernet noted in [5] that left-symmetric algebras feature in many fields, and Burde reviewed areas where left-symmetric algebras play key roles, including vector fields, rooted tree algebras, vertex algebras, operad theory, and affine manifolds (see [3]).

This paper begins by examining a structure analogous to Lie algebras, namely the Mock-Lie structure derived from antiassociative algebras, aiming to define Mock-Lie admissible algebras as associative counterparts of Lie admissible algebras. Next, we study the structure analogous to left-symmetric and refer to it as the quasi-antiassociative structure, defined via the antiassociator of the associated bilinear product.

The goal is to define Mock-Lie admissible algebras and investigate quasi-antiassociative algebras, with particular emphasis on their bimodules, matched pairs and bialgebras.

Since quasi-antiassociative structures differ from left-symmetric structures only by a sign (see Definition 2.5), we conjecture that the quasi-antiassociative structure may also correspond to a geometric connection.

The paper is organized as follows. Section 2 reviews basic concepts and properties of antiassociative and Mock-Lie algebras, and defines Mock-Lie admissible and quasi-antiassociative algebras. Section 3 covers bimodules and matched pairs of quasi-antiassociative algebras. Section 4 studies the quasi-antiassociative bialgebras. Section 5 concludes with final remarks.

2. Definitions and main properties

Throughout this work, we consider $\mathcal{A}$ a finite dimensional vector space over the field $\mathbb{K}$ of characteristic 0 together with a bilinear product " $\cdot$ " defined as : $\mathcal{A} \times \mathcal{A} \to \mathcal{A}$ such that $(x, y) \mapsto x \cdot y$.

Definition 2.1. [13] Let "$\cdot$" be a bilinear product in a vector space $\mathcal{A}$. Suppose that it satisfies the following law:

$$(x \cdot y) \cdot z = -x \cdot (y \cdot z).$$

Then, we call the pair $(\mathcal{A}, \cdot)$ an *antiassociative algebra.*

Definition 2.2. [15] An algebra $(\mathcal{A}, \diamond)$ over K is called *mock-Lie* if it is commutative $x \diamond y = y \diamond x$ and satisfies the Jacobi identity:

$$(x \diamond y) \diamond z + (z \diamond x) \diamond y + (y \diamond z) \diamond x = 0$$

for any x,y,$z \in \mathcal{A}$.

Theorem 2.3. [15] *Given an antiassociative algebra $(\mathcal{A}, \cdot)$, the new algebra $\mathcal{A}^\dagger$ with multiplication give by the "anticommutator"*

$$a \diamond b = (a \cdot b + b \cdot a)\,,$$

is a mock-Lie algebra.

Definition 2.4. A *Mock-Lie admissible algebra* is a nonassociative algebra which anticommutator is a Mock-Lie algebra.

Any antiassociative algebra is a Mock-Lie admissible algebra.

Definition 2.5. $(\mathcal{A}, \cdot)$, or simply $\mathcal{A}$, is said to be a *left quasi-antiassociative algebra* if $\forall\, x, y, z \in \mathcal{A}$, the antiassociator of the bilinear product $\cdot$ defined by $(x, y, z)_{-1} := (x \cdot y) \cdot z + x \cdot (y \cdot z)$, is symmetric in x and y, i.e.

$$(x, y, z)_{-1} = -(y, x, z)_{-1}.$$

As matter of notation simplification, we will denote $x \cdot y$ by xy if not any confusion.

Proposition 2.6. *Any antiassociative algebra is a left quasi-antiassociative algebra, and*

$$\mu \circ (\mu \otimes id) + \mu \circ (id \otimes \mu) + \mu \circ ((\mu \circ \tau) \otimes id) + \mu(id \otimes (\mu \circ \tau)) = 0,$$

where μ is the multiplication operator ant τ is such that $\tau(x \otimes y) = y \otimes x$.

Proof. Let $(\mathcal{A}, \cdot)$ be an antiassociative algebra. We have $\forall x, y, z \in \mathcal{A}$

$$(x \cdot y) \cdot z = -x \cdot (y \cdot z) \Leftrightarrow (x, y, z)_{-1} = 0 = -(y, x, z)_{-1}.$$

Thus $(x, y, z)_{-1} + (y, x, z)_{-1} = 0$, which implies that

$$(xy)z + x(yz) + (yx)z + y(xz) = 0.$$

Supposing μ be the bilinear product on $\mathcal{A}$, then the previous relation can be written as

$$\mu(\mu(x \otimes y) \otimes z) + \mu(x \otimes \mu(y \otimes z)) + \mu((\mu \circ \tau)(x \otimes y) \otimes z) + \mu(y \otimes (\mu \circ \tau) z \otimes x) = 0$$
$$\Leftrightarrow \mu \circ (\mu \otimes id) + \mu \circ (id \otimes \mu) + \mu \circ ((\mu \circ \tau) \otimes id) + \mu(id \otimes (\mu \circ \tau)) = 0. \quad \square$$

Definition 2.7. An algebra $(\mathcal{A}, \cdot)$ over $\mathbb{K}$ with the bilinear product given by $(x, y) \mapsto x \cdot y$ is called *right quasi-antiassociative* if the associator associated to the bilinear product on $\mathcal{A}$ is symmetric in right, i.e. for all $x, y, z \in \mathcal{A}$

$$(x, y, z)_{-1} = -(x, z, y)_{-1}.$$

Theorem 2.8. *The opposite algebra of the left quasi-antiassociative algebra is a right quasi-antiassociative algebra. The opposite algbera of the right quasi-antiassociative algebra is a left quasi-antiassociative algebra.*

Proof. Suppose that $(\mathcal{A}, \cdot)$ is a left quasi-antiassociative algebra and $(\mathcal{A}, \circ)$ the opposite algebra of the algebra $\mathcal{A}$. For any $x, y, z \in \mathcal{A}$ we have:

$$\begin{aligned}(x, y, z)_{-1,\circ} &= (x \circ y) \circ z + x \circ (y \circ z) = z \cdot (y \cdot x) + (z \cdot y) \cdot x \\ &= (z, y, x)_{-1} = -(y, z, x)_{-1} = -y \cdot (z \cdot x) - (y \cdot z) \cdot x \\ &= -(x \circ z) \circ y - x \circ (z \circ y) = -(x, z, y)_{-1,\circ}.\end{aligned}$$

Therefore, for all $x, y, z \in \mathcal{A}$, $(x, y, z)_{-1,\circ} = -(x, z, y)_{-1,\circ}$. Conversely, if $(\mathcal{A}, \circ)$ is a right quasi-antiassociative algebra, then by the same way we easily show that the algebra $(\mathcal{A}, \cdot)$ is left quasi-antiassociative. □

Theorem 2.8 proves the equivalence between left quasi-antiassociative algebras and right quasi-antiassociative algebras. Thus, for the rest of this paper, without any further clarification, left quasi-antiassociative algebras are called quasi-antiassociative algebras.

Considering the representations of the left L and right R multiplication operations:

$$\begin{array}{rcl} L : \mathcal{A} & \longrightarrow & \mathfrak{gl}(\mathcal{A}) \\ x & \longmapsto & L_x : \begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{A} \\ y & \longmapsto & x \cdot y, \end{array} \end{array}$$

$$\begin{array}{rcl} R : \mathcal{A} & \longrightarrow & \mathfrak{gl}(\mathcal{A}) \\ x & \longmapsto & R_x : \begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{A} \\ y & \longmapsto & y \cdot x, \end{array} \end{array}$$

we infer the adjoint representation $\mathrm{ad} := L + R$ of the sub-adjacent Mock-Lie algebra of a quasi-antiassociative algebra $\mathcal{A}$ as follows:

$$\begin{array}{rcl} \mathrm{ad} : \mathcal{A} & \longrightarrow & \mathfrak{gl}(\mathcal{A}) \\ x & \longmapsto & \mathrm{ad}_x : \begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{A} \\ y & \longmapsto & [x, y], \end{array} \end{array}$$

such that $\forall\, x, y \in \mathcal{A}, \mathrm{ad}_x(y) := (L_x + R_x)(y)$.

Definition 2.9. [15] A vector space V is a *module over a Mock-Lie algebra* $\mathcal{A}$, if there is a linear map (a representation) $\rho : \mathcal{A} \to End(V)$ such that

$$\rho(x \diamond y)(v) = -\rho(x)(\rho(y)v) - \rho(y)(\rho(x)v)$$

for any $x, y \in \mathcal{A}$ and $v \in V$.

Proposition 2.10. *Let $(\mathcal{A}, \cdot)$ be a quasi-antiassociative algebra. For any $x, y \in \mathcal{A}$, the following relations are satisfied:*

- *The anticommutator associated to the bilinear product " $\cdot$ " given by $[x, y] = x \cdot y + y \cdot x$ defines a Mock-Lie algebra structure on $\mathcal{A}$.*
- *The left multiplication operator gives a representation of the Mock-Lie algebra, that is*

  $$L_{[x,y]} = -[L_x, L_y],$$

  and the following relation is also satisfied

  $$[L_x, R_y] = -R_{x \cdot y} - R_y R_x,$$

  where the linear map R is the right multiplication operator associated to the bilinear product " $\cdot$ " on $\mathcal{A}$.
- $[L_x, R_y] = -[R_x, L_y]$
- $L_{xy} + L_x L_y = -L_{yx} - L_y L_x$,
- *$ad = L + R$ a linear representation of the sub-adjacent Mock-Lie algebra of $(\mathcal{A}, \cdot)$ and,*

  $$[ad_x, ad_y] = ad_{_{[x,y]}}.$$

Proof. Consider the quasi-antiassociative algebra $(\mathcal{A}, \cdot)$. For any $x, y, z \in \mathcal{A}$ we have

$$\begin{aligned}
&[x,[y,z]] + [y,[z,x]] + [z,[y,x]] = [x, yz + zy] + [y, zx + xz] + [z, xy + yx] \\
&= x(yz) + x(zy) + (yz)x + (zy)x + y(zx) + y(xz) + (zx)y + (xz)y + z(xy) \\
&\qquad + z(yx) + (xy)z + (yx)z \\
&= \{x(yz) + (yz)x + y(zx) + (zx)y + z(xy) + (xy)z\} \\
&\qquad + \{(zy)x + x(zy) + y(xz) + (xz)y + z(yx) + (yx)z\} \\
&= \{(x,y,z)_{_{-1}} + (y,z,x)_{_{-1}} + (z,x,y)_{_{-1}}\} \\
&\qquad + \{(z,y,x)_{_{-1}} + (x,z,y)_{_{-1}} + (y,x,z)_{_{-1}}\} \\
&= \{(y,x,z)_{_{-1}} + (x,y,z)_{_{-1}}\} + \{(z,y,x)_{_{-1}} + (y,z,x)_{_{-1}}\} + \{(x,z,y)_{_{-1}}
\end{aligned}$$

$$+(z,x,y)_{-1}\} = 0.$$

On other hand, we have for all $x, y, z \in \mathcal{A}$

$$\begin{aligned}(x,y,z)_{-1} = -(y,x,z)_{-1} &\Leftrightarrow (xy)z + x(yz) = -(yx)z - y(xz)\\ &\Leftrightarrow (xy)z + (yx)z = -x(yz) - y(xz)\\ &\Leftrightarrow (L_{xy} + L_{yx})(z) = -(L_xL_y + L_yL_x)(z)\\ &\Leftrightarrow (L_{xy} + L_{yx})(z) = -[L_x, L_y](z)\end{aligned}$$

Therefore, $L_{[x,y]} = -[L_x, L_y]$ for all $x, y \in \mathcal{A}$ holds.

We have for all $x, y, z \in \mathcal{A}$

$$\begin{aligned}(x,y,z)_{-1} = -(y,x,z)_{-1} &\Leftrightarrow (xy)z + x(yz) = -(yx)z - y(xz)\\ &\Leftrightarrow (R_zL_x - L_xR_z)(y) = -(R_zR_x + R_{xz})(y)\\ &\Leftrightarrow [R_z, L_x](y) = -(R_zR_x + R_{xz})(y),\end{aligned}$$

Therefore, the following relation holds

$$[L_x, R_y] = -(R_{xy} + R_yR_x).$$

We have for all $x, y, z \in \mathcal{A}$

$$\begin{aligned}[L_x, R_y](z) &= L_x\left(R_y(z)\right) + R_y\left(L_x(z)\right)\\ &= x(zy) + (xz)y = (x,z,y)_{-1} = -(z,x,y)_{-1}\\ &= -\left((zx)y + z(xy)\right) = -\left(R_y\left(R_x(z)\right) + R_{xy}(z)\right)\\ &= -\left(R_yR_x + R_{xy}\right)(z)\\ &= (z,y,x)_{-1} = (zy)x + z(yx) = R_xR_y(z) + R_{xy}(z)\\ &= -(y,z,x) - (yz)x - y(zx) = -R_xL_y(z) - L_yR_x(z)\\ &= -[R_x, L_y](z).\end{aligned}$$

Therefore $[L_x, R_y] = -[R_x, L_y]$.

We have for all $x, y, z \in \mathcal{A}$

$$\begin{aligned}\left(L_{xy} + L_xL_y\right)(z) &= L_{xy}(z) + L_x\left(L_y(z)\right) = (xy)z + x(yz)\\ &= (x,y,z)_{-1} = -(y,x,z)_{-1} = -\left((yx)z + y(xz)\right)\\ &= -\left(L_{yx}(z) + L_y(L_x(z))\right)\\ &= -\left(L_{yx} + L_yL_x\right)(z).\end{aligned}$$

Thus $L_{xy} + L_xL_y = -L_{yx} - L_yL_x$.

We have for all $x, y \in \mathcal{A}$,

$$\begin{aligned}[ad_x, ad_y] &= [L_x, L_y] + [L_x, R_y] + [R_x, L_y] + [R_x, R_y] \\ &= [L_x, L_y] + [R_x, R_y] \\ &= (L_x L_y + R_x R_y) + (L_y L_x + R_y R_x) \\ &= (L + R)_{xy} + (L + R)_{yx} = (L + R)_{xy+yx} \\ &= (L + R)_{[x,y]} = ad_{[x,y]}\end{aligned}$$

This completes the proof. □

Remark 2.11. From this proposition, quasi-antiassociative algebras are Mock-Lie admissible algebras.

Thus, $(\mathcal{A}, \cdot)$ can be called the compatible quasi-antiassociative algebra product of the Mock-Lie algebra $\mathcal{G}(\mathcal{A})$.

3. Bimodules and matched pairs

Definition 3.1. Let $\mathcal{A}$ be a quasi-antiassociative algebra, V be a vector space. Suppose $l, r : \mathcal{A} \to \mathfrak{gl}(V)$ be two linear maps such that

$$[l_x, r_y] = -[l_y, r_x] \tag{3.1}$$

$$l_{xy} + l_x l_y = -l_{yx} - l_y l_x \tag{3.2}$$

for all $x, y \in \mathcal{A}$. Then, (l, r, V) (or simply (l, r)) is called *bimodule of the quasi-antiassociative algebra* $\mathcal{A}$.

Proposition 3.2. *Let $(\mathcal{A}, \cdot)$ be a quasi-antiassociative algebra and V be a vector space over $\mathbb{K}$. Consider two linear maps, $l, r : \mathcal{A} \to \mathfrak{gl}(V)$. Then, (l, r, V) is a bimodule of $\mathcal{A}$ if and only if, the semidirect sum $\mathcal{A} \oplus V$ of vector spaces is turned into a quasi-antiassociative algebra by defining the multiplication in $\mathcal{A} \oplus V$ by $\forall x_1, x_2 \in \mathcal{A},\ v_1, v_2 \in V$,*

$$(x_1 + v_1) * (x_2 + v_2) = x_1 \cdot x_2 + (l_{x_1} v_2 + r_{x_2} v_1).$$

We denote it by $\mathcal{A} \ltimes_{l,r}^{-1} V$ or simply $\mathcal{A} \ltimes^{-1} V$.

Proof. It is obvious that the semidirect sum of two vector spaces is also a vector space. Now suppose that (l, r, V) is a bimodule of $\mathcal{A}$ and show that $(\mathcal{A} \oplus V, *)$ is a quasi-antiassociative algebra. Since $*$ is a bilinear product, for all $x_1, x_2, x_3 \in \mathcal{A}$ and for all $v_1, v_2, v_3 \in V$, we have:

$$
\begin{aligned}
&((x_1+v_1),(x_2+v_2),(x_3+v_3))_{-1}\\
&\quad=\{(x_1+v_1)*(x_2+v_2)\}*(x_3+v_3)+(x_1+v_1)*\{(x_2+v_2)*(x_3+v_3)\}\\
&\quad=(x_1x_2)x_3+l_{x_1x_2}v_3+r_{x_3}(l_{x_1}v_2+r_{x_2}v_1)+x_1(x_2x_3)+l_{x_1}(l_{x_2}v_3)\\
&\qquad\qquad+l_{x_1}(r_{x_3}v_2)+r_{x_2x_3}v_1\\
&\quad=(x_1x_2)x_3+l_{x_1x_2}v_3+r_{x_3}(l_{x_3}v_2)+r_{x_3}(r_{x_2}v_1)+x_1(x_2x_3)+l_{x_1}(l_{x_2}v_3)\\
&\qquad\qquad+l_{x_1}(r_{x_3}v_2)+r_{x_2x_3}v_1,\\
&((x_1+v_1),(x_2+v_2),(x_3+v_3))\\
&\quad=(x_1,x_2,x_3)_{-1}+(l_{x_1x_2}+l_{x_1}l_{x_2})v_3+(r_{x_3}l_{x_1}+l_{x_1}r_{x_3})v_2+(r_{x_3}r_{x_2}+r_{x_2x_3})v_1.\\
&((x_2+v_2),(x_1+v_1),(x_3+v_3))\\
&\quad=(x_2,x_1,x_3)+(l_{x_2x_1}+l_{x_2}l_{x_1})v_3+(r_{x_3}l_{x_2}+l_{x_2}r_{x_3})v_1+(r_{x_3}r_{x_1}+r_{x_1x_3})v_2.
\end{aligned}
$$

Therefore,

$$
\begin{aligned}
(x_1+v_1,x_2+v_2,x_3+v_3)_{-1}&=-(x_3+v_3,x_2+v_2,x_1+v_1)_{-1}\\
&\Leftrightarrow\begin{cases} l_{x_1x_2}+l_{x_1}l_{x_2}=-l_{x_2x_1}-l_{x_2}l_{x_1}\\ r_{x_3}l_{x_1}+l_{x_1}r_{x_3}=-r_{x_3}r_{x_1}-r_{x_1x_3}\\ r_{x_2x_3}+r_{x_3}r_{x_2}=-r_{x_3}l_{x_2}-l_{x_2}r_{x_3}\end{cases}\\
&\Leftrightarrow(\mathcal{A}\oplus V,*)\text{ is a quasi-antiassociative algebra.}\qquad\square
\end{aligned}
$$

Furthermore, we derive the next result.

Proposition 3.3. *Let $\mathcal{A}$ be a quasi-antiassociative algebra and V be a vector space over $\mathbb{K}$. Consider two linear maps, $l,r:\mathcal{A}\to\mathfrak{gl}(V)$, such that (l,r,V) is a bimodule of $\mathcal{A}$. Then, the map: $l+r:\mathcal{A}\longrightarrow\mathfrak{gl}(V)$ $x\longmapsto l_x+r_x$, is a linear representation of the sub-adjacent Mock-Lie algebra of $\mathcal{A}$.*

Proof. Let (l,r,V) be a bimodule of the quasi-antiassociative algebra $\mathcal{A}$. Then, $\forall x,y\in\mathcal{A}\quad[l_x,r_y]=-[l_y,r_x];l_{xy}+l_xl_y=-l_yl_x-l_{yx}$. Besides, it is a matter of straightforward computation to show that $l+r$ is a linear map on $\mathcal{A}$. Then, we have:

$$
\begin{aligned}
[(l+r)(x),(l+r)(y)]&=[l_x+r_x,l_y+r_y]=[l_x,l_y]+[l_x,r_y]+[r_x,l_y]+[r_x,r_y]\\
&=[l_x,l_y]+[r_x,r_y]\\
&=l_xl_y+l_yl_x+r_xr_y+r_yr_x\\
&=\{l_xl_y+r_xr_y\}+\{l_yl_x+r_yr_x\}\\
&=\{l_{xy}+r_{xy}\}+\{l_{yx}+r_{yx}\}=(l+r)_{xy}+(l+r)_{yx}\\
&=(l+r)_{[x,y]}.
\end{aligned}
$$

Therefore, (l,r,V) is a bimodule of $\mathcal{A}$ implies that $l+r$ is a representation of the linear representation of the sub-adjacent Mock-Lie algebra of $\mathcal{A}$. $\square$

Example 3.4. *According to the Proposition 2.10, one can deduce that* $(L, R, \mathcal{A})$ *is a bimodule of the quasi-antiassociative algebra* $\mathcal{A}$*, where* L *and* R *are the left and right multiplication operator representations, respectively.*

Definition 3.5. Let $\mathcal{G}$ and $\mathcal{H}$ be two Mock-Lie algebras and let $\mu : \mathcal{H} \to \mathfrak{gl}(\mathcal{G})$ and $\rho : \mathcal{G} \to \mathfrak{gl}(\mathcal{H})$ be two Mock-Lie algebra representations. Then, $(\mathcal{G}, \mathcal{H}, \rho, \mu)$ is called a *matched pair* of the Mock-Lie algebras $\mathcal{G}$ and $\mathcal{H}$, denoted by $\mathcal{H} \bowtie_{\mu}^{-1,\rho} \mathcal{G}$ if and only if μ and ρ satisfy: for all $x, y \in \mathcal{G}, a, b \in \mathcal{H}$,

$$\rho(x)\,[a,b] + [\rho(x)a, b] + [a, \rho(x)b] + \rho(\mu(a)x)b + \rho(\mu(b)x)a = 0, \tag{3.3}$$

$$\mu(a)\,[x,y] + [\mu(a)x, y] + [x, \mu(a)y] + \mu(\rho(x)a)y + \mu(\rho(y)a)x = 0. \tag{3.4}$$

In this case, $(\mathcal{G} \oplus \mathcal{H}, [,])$ defines a Mock-Lie algebra with respect to the product $*$ satisfying:

$$[(x+a), (y+b)] = [x,y] + \mu(a)y + \mu(b)x + [a,b] + \rho(x)b + \rho(y)a.$$

Theorem 3.6. *Let* $(\mathcal{A}, \cdot)$ *and* $(\mathcal{B}, \circ)$ *be two quasi-antiassociative algebras. Suppose that* $(l_{\mathcal{A}}, r_{\mathcal{A}}, \mathcal{B})$ *and* $(l_{\mathcal{B}}, r_{\mathcal{B}}, \mathcal{A})$ *are bimodules of* $\mathcal{A}$ *and* $\mathcal{B}$*, respectively, obeying the relations:*

$$r_{\mathcal{A}}(x)([a,b]) = r_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a + r_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b + a \circ (r_{\mathcal{A}}(x)b) + b \circ (r_{\mathcal{A}}(x)a), \tag{3.5}$$

$$\begin{aligned} -l_{\mathcal{A}}(x)(a \circ b) &= l_{\mathcal{A}}(l_{\mathcal{B}}(a)x + r_{\mathcal{B}}(a)x)b + (l_{\mathcal{A}}(x)a + r_{\mathcal{A}}(x)a) \circ b \\ &+ r_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a + a \circ (l_{\mathcal{A}}(x)b), \end{aligned} \tag{3.6}$$

$$r_{\mathcal{A}}(a)[x,y] = r_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x + r_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y + x(r_{\mathcal{B}}(a)y) + y(r_{\mathcal{B}}(a)x), \tag{3.7}$$

$$\begin{aligned} -l_{\mathcal{B}}(a)(xy) &= l_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y + (r_{\mathcal{B}}(a)x)y + x(l_{\mathcal{B}}(a)y) + r_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x \\ &+ (l_{\mathcal{B}}(a)x)y + l_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y, \end{aligned} \tag{3.8}$$

for all $x, y \in A$ *and* $a, b \in \mathcal{B}$*. Then, there is a quasi-antiassociative algebra structure on* $\mathcal{A} \oplus \mathcal{B}$ *given by:*

$$(x+a) * (y+b) = (x \cdot y + l_{\mathcal{B}}(a)y + r_{\mathcal{B}}(b)x) + (a \circ b + l_{\mathcal{A}}(x)b + r_{\mathcal{A}}(y)a). \tag{3.9}$$

We denote this quasi-antiassociative algebra by $\mathcal{A} \bowtie_{l_{\mathcal{B}}, r_{\mathcal{B}}}^{-1, l_{\mathcal{A}}, r_{\mathcal{A}}} \mathcal{B}$, or simply by $\mathcal{A} \bowtie^{-1} \mathcal{B}$. Then $(\mathcal{A}, \mathcal{B}, l_{\mathcal{A}}, r_{\mathcal{A}}, l_{\mathcal{B}}, r_{\mathcal{B}})$ satisfying the above conditions is called matched pair of the quasi-antiassociative algebras $\mathcal{A}$ and $\mathcal{B}$.

Proof. Consider $x, y \in \mathcal{A}$ and $a, b \in \mathcal{B}$. We have

$$\begin{aligned} (x+a) * (y+b) &= (xy + l_{\mathcal{B}}(a)y + r_{\mathcal{B}}(b)x) + (a \circ b + l_{\mathcal{A}}(x)b + r_{\mathcal{A}}(y)a), \\ x * a &= r_{\mathcal{B}}(a)x + l_{\mathcal{A}}(x)a, \ b = 0, y = 0, \\ a * y &= l_{\mathcal{B}}(a)x + r_{\mathcal{A}}(y)a, \ x = 0, b = 0. \end{aligned}$$

Since $(x,y,z)_{-1} = (xy)z + x(yz)$, we have

$$\begin{aligned}(x,b,c)_{-1} &= (x*b)*c + x*(b\circ c)(r_{\mathcal{B}}(b)x + l_{\mathcal{A}}(x)x)*c + x*(b\circ c)\\ &= r_{\mathcal{B}}(c)(r_{\mathcal{B}}(b)x) + (l_{\mathcal{A}}(x)b)\circ c + l_{\mathcal{A}}(r_{\mathcal{B}}(b)x)c + r_{\mathcal{B}}(b\circ c)x + l_{\mathcal{A}}(x)(b\circ c),\end{aligned}$$

$$\begin{aligned}(a,y,c)_{-1} &= (a*y)*c + a*(y*c)\\ &= (l_{\mathcal{B}}(a)y + r_{\mathcal{A}}(y)a)*c + a*(l_{\mathcal{A}}(y)c + r_{\mathcal{B}}(c)y)\\ &= l_{\mathcal{A}}(l_{\mathcal{B}}(a)y)c + r_{\mathcal{B}}(c)(l_{\mathcal{B}}(a)y) + (r_{\mathcal{A}}(y)a)\circ c\\ &+ a\circ(l_{\mathcal{A}}(y)c) + l_{\mathcal{B}}(a)(r_{\mathcal{B}}(c)y) + r_{\mathcal{A}}(r_{\mathcal{B}}(c)y)a,\end{aligned}$$

$$\begin{aligned}(x,y,c)_{-1} &= (x\cdot y)*c + x*(y*c)\\ &= (l_{\mathcal{A}}(x\cdot y)c + r_{\mathcal{B}}(c)(x\cdot y)*c + x*(l_{\mathcal{A}}(y)c + r_{\mathcal{B}}(c)y)\\ &= l_{\mathcal{A}}(xy)c + r_{\mathcal{B}}(c)(xy) + x\cdot(r_{\mathcal{B}}(c)y) + l_{\mathcal{A}}(x)(l_{\mathcal{A}}(y)c) + r_{\mathcal{B}}(l_{\mathcal{A}}(y)c)x,\end{aligned}$$

$$\begin{aligned}(x,b,z)_{-1} &= (x*b)*z + x*(b*z)\\ &= (l_{\mathcal{A}}(x)b + r_{\mathcal{B}}(b)x)*z + x*(l_{\mathcal{B}}(b)z + r_{\mathcal{A}}(z)b)\\ &= l_{\mathcal{B}}(l_{\mathcal{A}}(x)b)z + r_{\mathcal{A}}(z)(l_{\mathcal{A}}(x)b) + (r_{\mathcal{B}}(b)x)\cdot z\\ &+ x\cdot(l_{\mathcal{B}}(b)z) + l_{\mathcal{A}}(x)(r_{\mathcal{A}}(z)b) + r_{\mathcal{B}}(r_{\mathcal{A}}(z)b)x,\end{aligned}$$

$$(a,b,c)_{-1} = (a\circ b)\circ c + a\circ(b\circ c)$$

$$\begin{aligned}(a,y,z)_{-1} &= (a*y)*z + a*(y\cdot z)\\ &= (l_{\mathcal{B}}(a)y + r_{\mathcal{A}}(y)a)*z + l_{\mathcal{B}}(a)(y\cdot z) + r_{\mathcal{A}}(y\cdot z)a\\ &= (l_{\mathcal{B}}(a)y)z + l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)z + r_{\mathcal{A}}(z)(r_{\mathcal{A}}(y)a) + l_{\mathcal{B}}(a)(y\cdot z) + r_{\mathcal{A}}(y\cdot z)a,\end{aligned}$$

$$\begin{aligned}(a,b,z)_{-1} &= (a\circ b)*z + a*(b*z)\\ &= (l_{\mathcal{B}}(a\circ a)\cdot z + r_{\mathcal{A}}(z)(a\circ b)) + a*(l_{\mathcal{B}}(b)z + r_{\mathcal{A}}(z)b)\\ &= l_{\mathcal{B}}(a\circ b)z + r_{\mathcal{A}}(z)(\circ b) + a\circ(r_{\mathcal{A}}(z)b) + l_{\mathcal{B}}(a)(l_{\mathcal{B}}(b)z) - r_{\mathcal{A}}(l_{\mathcal{B}}(b)z)a,\end{aligned}$$

The first part of the associator reads:

$$\begin{aligned}&\{(x+a)*(y+b)\}*(z+c)\\ &= \{(xy + l_{\mathcal{B}}(a)y + r_{\mathcal{B}}(b)x) + (a\circ b + l_{\mathcal{A}}(y) + r_{\mathcal{A}}(y)a)\}*(z+c)\\ &= (xy + l_{\mathcal{B}}(a)y + r_{\mathcal{B}}(b)x)z + l_{\mathcal{B}}(a\circ b + l_{\mathcal{A}}(x)b + r_{\mathcal{A}}(y)a)z\\ &\quad + r_{\mathcal{B}}(c)(xy + l_{\mathcal{B}}(a)y + r_{\mathcal{B}}(b)x) + (a\circ b + l_{\mathcal{A}}(x)b + r_{\mathcal{A}}(y)a)\circ c\\ &\quad + l_{\mathcal{A}}(a)(yz + l_{\mathcal{B}}(b)z + r_{\mathcal{B}}(c)y) + r_{\mathcal{A}}(z)(a\circ bl_{\mathcal{A}}(x)b + r_{\mathcal{A}}(y)a)\\ &= (xy)z + (l_{\mathcal{B}}(a)y)z + (r_{\mathcal{B}}(b)x)z + l_{\mathcal{B}}(a\circ b)z + l_{\mathcal{B}}(l_{\mathcal{A}}(x)b)z + l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)z\\ &\quad + r_{\mathcal{B}}(c)(xy) + r_{\mathcal{B}}(c)(l_{\mathcal{B}}(a)y) + r_{\mathcal{B}}(c)(r_{\mathcal{B}}(b)x)) + (a\circ b)\circ c + (l_{\mathcal{A}}(x)b)\circ c\\ &\quad + (r_{\mathcal{A}}(y)a)\circ c + l_{\mathcal{A}}(a)(yz) + l_{\mathcal{A}}(a)(l_{\mathcal{B}}(b)z) + l_{\mathcal{A}}(a)(r_{\mathcal{B}}(c)y)\\ &\quad + r_{\mathcal{A}}(z)(a\circ b) + r_{\mathcal{A}}(z)(l_{\mathcal{A}}(x)b) + r_{\mathcal{A}}(z)(r_{\mathcal{A}}(y)a)\end{aligned}$$

while its second part:

$$(x+a)*\{(y+b)*(z+c)\}$$

$$
\begin{aligned}
&= (x+a) * \{yz + l_{\mathcal{B}}(b)z + r_{\mathcal{B}}(c)y + b \circ c + l_{\mathcal{A}}(y)c + r_{\mathcal{A}}(z)b\} \\
&= x(yz + l_{\mathcal{B}}(b)z + r_{\mathcal{B}}(c)y) + l_{\mathcal{B}}(a)(yz + l_{\mathcal{B}}(b)z + r_{\mathcal{B}}(c)y) \\
&\quad + r_{\mathcal{B}}(b \circ c + l_{\mathcal{A}}(y)c + r_{\mathcal{A}}(z)b)x + a \circ (b \circ c + l_{\mathcal{A}}(y)c \\
&\quad + r_{\mathcal{A}}(z)b) + l_{\mathcal{A}}(x)(b \circ c + l_{\mathcal{A}}(y)c + r_{\mathcal{A}}(z)b) + r_{\mathcal{A}}(yz + l_{\mathcal{B}}(b)z + r_{\mathcal{B}}(c)y)a \\
&= x(yz) + x(l_{\mathcal{B}}(b)z) + x(r_{\mathcal{B}}(c)y) + l_{\mathcal{B}}(a)(yz) + l_{\mathcal{B}}(a)(l_{\mathcal{B}}(b)z) \\
&\quad + l_{\mathcal{B}}(a)(r_{\mathcal{B}}(c)y) + r_{\mathcal{B}}(b \circ c)x + r_{\mathcal{B}}(l_{\mathcal{A}}(y)c)x + r_{\mathcal{B}}(r_{\mathcal{A}}(z)b)x \\
&\quad + a \circ (b \circ c) + a \circ (l_{\mathcal{A}}(y)c) + a \circ (r_{\mathcal{A}}(z)b) + l_{\mathcal{A}}(x)(b \circ c) \\
&\quad + l_{\mathcal{A}}(x)(l_{\mathcal{A}}(y)c) + l_{\mathcal{A}}(x)(r_{\mathcal{A}}(z)b) + r_{\mathcal{A}}(yz)a + r_{\mathcal{A}}(l_{\mathcal{B}}(b)z)a + r_{\mathcal{A}}(r_{\mathcal{B}}(c)y)a
\end{aligned}
$$

and the associator takes the form:

$$
\begin{aligned}
&(x+a, y+b, z+c)_{-1} \\
&= \{(xy)z - x(yz)\} + \{(a \circ b) \circ c - a \circ (b \circ c)\} \\
&\quad + \{r_{\mathcal{B}}(c)(r_{\mathcal{B}}(b)x) + (l_{\mathcal{A}}(x)b) \circ c + l_{\mathcal{A}}(r_{\mathcal{B}}(b)x)c \\
&\quad - r_{\mathcal{B}}(b \circ c)x - l_{\mathcal{A}}(x)(b \circ c)\}\{l_{\mathcal{A}}(l_{\mathcal{B}}()a)yc + r_{\mathcal{B}}(c)(l_{\mathcal{B}}(a)y) + (r_{\mathcal{A}}y) \circ c \\
&\quad - a \circ (l_{\mathcal{A}}(y)c) - l_{\mathcal{B}}(a)(r_{\mathcal{B}}(c)y) - r_{\mathcal{A}}(r_{\mathcal{B}}(c)y)a\} + \{l_{\mathcal{A}}(l_{\mathcal{B}}(a)y)c \\
&\quad + r_{\mathcal{B}}(c)(l_{\mathcal{B}}(a)y) + (r_{\mathcal{A}}(y)a) \circ c - a \circ (l_{\mathcal{A}}(y)c) - l_{\mathcal{B}}(a)(r_{\mathcal{B}}(c)y) \\
&\quad - r_{\mathcal{A}}(r_{\mathcal{B}}(c)y)a\} + \{l_{\mathcal{B}}(l_{\mathcal{A}}(x)b)z + r_{\mathcal{A}}(z)(l_{\mathcal{A}}(x)b) + (r_{\mathcal{B}}(b)x)z \\
&\quad - x(l_{\mathcal{B}}(b)z) - l_{\mathcal{A}}(x)(r_{\mathcal{A}}(z)b)x\} + \{(l_{\mathcal{B}}(a)y)z + l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)z \\
&\quad + r_{\mathcal{A}}(z)(r_{\mathcal{A}}(y)a) - l_{\mathcal{B}}(a)(yz) - r_{\mathcal{A}}(yz)a\} + \{l_{\mathcal{B}}(a \circ b)z \\
&\quad + r_{\mathcal{A}}(z)(a \circ b) - a \circ (r_{\mathcal{A}}(z)b) - l_{\mathcal{B}}(a)(l_{\mathcal{B}}(b)z) - r_{\mathcal{A}}(l_{\mathcal{B}}(b)z)a\} \\
&= (x, y, z)_{-1} + (a, b, c)_{-1} + \{r_{\mathcal{B}}(c)(x \cdot y) + l_{\mathcal{A}}(x \cdot y)c + x \cdot (r_{\mathcal{B}}(c)y) \\
&\quad + l_{\mathcal{A}}(x)(l_{\mathcal{A}}(y)c) + r_{\mathcal{B}}(l_{\mathcal{A}}(y)c)x\} + \{r_{\mathcal{B}}(c)(r_{\mathcal{B}}(b)x) + l_{\mathcal{A}}(r_{\mathcal{B}}(b)x)c \\
&\quad + r_{\mathcal{B}}(b \circ c)x + (l_{\mathcal{A}}(x)b) \circ c + l_{\mathcal{A}}(x)(b \circ c)\} + \{(r_{\mathcal{B}}(b)x) \cdot z + l_{\mathcal{B}}(l_{\mathcal{A}}(x)b)z \\
&\quad + r_{\mathcal{A}}(z)(l_{\mathcal{A}}(x)b) + x \cdot (l_{\mathcal{B}}(b)z) + r_{\mathcal{B}}(r_{\mathcal{A}}(z)b)x + l_{\mathcal{A}}(x)(r_{\mathcal{A}}(z)b)\} \\
&\quad + \{(l_{\mathcal{B}}(a)y) \cdot z + l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)z + r_{\mathcal{A}}(z)(r_{\mathcal{A}}(y)a) + l_{\mathcal{B}}(a)(y \cdot z) \\
&\quad + r_{\mathcal{A}}(y \cdot z)a\} + \{r_{\mathcal{B}}(c)(l_{\mathcal{B}}(a)y) + (r_{\mathcal{A}}(y)a) \circ c + l_{\mathcal{A}}(l_{\mathcal{B}}(a)y)c \\
&\quad + l_{\mathcal{B}}(a)(r_{\mathcal{B}}(c)y) + a \circ (l_{\mathcal{A}}(y)c) + r_{\mathcal{A}}(r_{\mathcal{B}}(c)y)a\} + \{l_{\mathcal{B}}(a \circ b)z \\
&\quad + r_{\mathcal{A}}(z)(a \circ b) + l_{\mathcal{B}}(a)(l_{\mathcal{B}}(b)z) + a \circ (r_{\mathcal{A}}(z)b) + r_{\mathcal{A}}(l_{\mathcal{B}}(b)z)a\}.
\end{aligned}
$$

Further, we have

$$
\begin{aligned}
(x, y, c)_{-1} &= r_{\mathcal{B}}(c)(x \cdot y) + x \cdot (r_{\mathcal{B}}(c)y) + r_{\mathcal{B}}(l_{\mathcal{A}}(y)c)x + l_{\mathcal{A}}(x \cdot y)c + l_{\mathcal{A}}(x)(l_{\mathcal{A}}(y)c), \\
(x, b, c)_{-1} &= r_{\mathcal{B}}(c)(r_{\mathcal{B}}(b)x) + l_{\mathcal{A}}(r_{\mathcal{B}}(b)x)c + (l_{\mathcal{A}}(x)b) \circ c + r_{\mathcal{B}}(b \circ c)x + l_{\mathcal{A}}(x)(b \circ c), \\
(x, b, z)_{-1} &= (r_{\mathcal{B}}(b)x) \cdot z + l_{\mathcal{B}}(l_{\mathcal{A}}(x)b)z + r_{\mathcal{A}}(z)(l_{\mathcal{A}}(x)b) + x \cdot (l_{\mathcal{B}}(b)z) \\
&\quad + r_{\mathcal{B}}(r_{\mathcal{A}}(z)b)x + l_{\mathcal{A}}(x)(r_{\mathcal{A}}(z)b), \\
(a, y, z)_{-1} &= (l_{\mathcal{B}}(a)y) \cdot z + l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)z + r_{\mathcal{A}}(z)(r_{\mathcal{A}}(y)a) + l_{\mathcal{B}}(a)(y \cdot z) + r_{\mathcal{A}}(y \cdot z)a, \\
(a, y, c)_{-1} &= r_{\mathcal{B}}(c)(l_{\mathcal{B}}(a)y) + (r_{\mathcal{A}}(y)a) \circ c + l_{\mathcal{A}}(l_{\mathcal{B}}(a)y)c + l_{\mathcal{B}}(a)(r_{\mathcal{B}}(c)y) \\
&\quad + a \circ (l_{\mathcal{A}}(y)c) + r_{\mathcal{A}}(r_{\mathcal{B}}(c)y)a, \\
(a, b, z)_{-1} &= l_{\mathcal{B}}(a \circ b)z + r_{\mathcal{A}}(z)(a \circ b) + l_{\mathcal{B}}(a)(l_{\mathcal{B}}(b)z) + r_{\mathcal{A}}(l_{\mathcal{B}}(b)z)a + r_{\mathcal{A}}(l_{\mathcal{B}}(b)z)a \\
&\quad + a \circ (r_{\mathcal{A}}(z)b),
\end{aligned}
$$

which can also be re-expressed as:

$$(x+a,y+b,z+c)_{-1} = (x,y,z)_{-1} + (x,y,c)_{-1} + (x,b,z)_{-1} + (x,b,c)_{-1} \\ +(a,y,z)_{-1} + (a,y,c)_{-1} + (a,b,z)_{-1} + (a,b,c)_{-1}.$$

Similarly,

$$(y+b,x+a,z+c)_{-1} = (y,x,z)_{-1} + (b,x,z)_{-1} + (y,x,c_{-1}) + (b,a,z)_{-1} \\ +(y,a,z)_{-1} + (y,a,c)_{-1} + (b,a,c)_{-1} + (b,x,c)_{-1}.$$

Since $\mathcal{A}$ and $\mathcal{B}$ are quasi-antiassociative algebras, we have

$$(x,y,z)_{-1} = -(y,x,z)_{-1} \\ (a,b,c)_{-1} = -(b,a,c)_{-1}.$$

Hence,

$$(x,b,z)_{-1} = -(b,x,z)_{-1} \Leftrightarrow (y,a,z)_{-1} = -(a,y,z)_{-1}\{x \to y, b \to a, z \to z\} \\ (x,b,c)_{-1} = -(b,x,c)_{-1} \Leftrightarrow (a,y,c)_{-1} = -(y,a,c)_{-1}\{x \to y, b \to a, c \to c\}.$$

Then, it remains to show that:

$$(x,a,y)_{-1} = -(a,x,y)_{-1}, \tag{3.10}$$
$$(x,a,b)_{-1} = -(a,x,b)_{-1}, \tag{3.11}$$
$$(x,y,a)_{-1} = -(y,x,a)_{-1}, \tag{3.12}$$
$$(a,b,x)_{-1} = -(b,a,x)_{-1}. \tag{3.13}$$

We have

$$\begin{aligned}(3.10) \Leftrightarrow\ & l_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y + r_{\mathcal{A}}(y)(l_{\mathcal{A}}(x)a) + (r_{\mathcal{B}}(a)x)y + x(l_{\mathcal{B}}(a)y) \\ & +l_{\mathcal{A}}(x)(r_{\mathcal{A}}(y)a) + r_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x \\ =\ & -\{(l_{\mathcal{B}}(a)x)y + l_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y + r_{\mathcal{A}}(y)(r_{\mathcal{A}}(x)a) \\ & +l_{\mathcal{B}}(a)(xy) + r_{\mathcal{A}}(xy)a\} \\ \Leftrightarrow\ & -l_{\mathcal{B}}(a)(xy) = l_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y + (r_{\mathcal{B}}(a)x)y + x(l_{\mathcal{B}}(a)y) + \\ & +r_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x + (l_{\mathcal{B}}(a)x)y + l_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y + l_{\mathcal{B}}(a)(xy)\end{aligned}$$

$(3.10) \Leftrightarrow (3.8)$ since$[l_x, r_y] = -r_{x\cdot y} - r_y r_x$.

$$\begin{aligned}(3.11) \Leftrightarrow\ & r_{\mathcal{B}}(b)(r_{\mathcal{B}}(a)x) + (l_{\mathcal{A}}(x)a) \circ b + l_{\mathcal{A}}(r_{\mathcal{B}}(a)x)b + r_{\mathcal{B}}(a \circ b)x \\ & +l_{\mathcal{A}}(x)(a \circ b) = -\{l_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b + r_{\mathcal{B}}(b)(l_{\mathcal{B}}(a)x) + (r_{\mathcal{A}}(x)a) \circ b \\ & +a \circ (l_{\mathcal{A}}(x)b) + l_{\mathcal{B}}(a)(r_{\mathcal{B}}(b)x) + r_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a\} \\ \Leftrightarrow\ & -l_{\mathcal{A}}(x)(a \circ b) = l_{\mathcal{A}}(l_{\mathcal{B}}(a)x + r_{\mathcal{B}}(a)x)b + (l_{\mathcal{A}}(x)a + r_{\mathcal{A}}(x)a) \circ b \\ & +r_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a + a \circ (l_{\mathcal{A}}(x)b) \text{ with, } (l_{\mathcal{B}}, r_{\mathcal{B}}) \text{ is bimodule of } \mathcal{B}\end{aligned}$$

$(3.11) \Leftrightarrow (3.6)$ with, $(l_{\mathcal{B}}, r_{\mathcal{B}})$ is bimodule of $\mathcal{B}$.

$$(3.12) \Leftrightarrow l_{\mathcal{A}}(xy)a + r_{\mathcal{B}}(a)(xy) + x(r_{\mathcal{B}}(a)y) + l_{\mathcal{A}}(x)(l_{\mathcal{A}}(y)a) + r_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x$$

$$= -\{l_{\mathcal{A}}(yx)a + r_{\mathcal{B}}(a)(yx) + y(r_{\mathcal{B}}(a)x) + l_{\mathcal{A}}(y)(l_{\mathcal{A}}(x)a) + r_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y\}$$
$$\Leftrightarrow r_{\mathcal{A}}(a)[x,y] = r_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x + r_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y + x(r_{\mathcal{B}}(a)y) + y(r_{\mathcal{B}}(a)x)$$

(3.12) $\Leftrightarrow$ (3.7), since (3.2) hold.

(3.13) $\Leftrightarrow l_{\mathcal{B}}(a\circ b)x + r_{\mathcal{A}}(x)(a\circ b) + a\circ(r_{\mathcal{A}}(x)b) + l_{\mathcal{B}}(a)(l_{\mathcal{B}}(b)x) + r_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a$
$= -\{l_{\mathcal{B}}(b\circ a)x + r_{\mathcal{A}}(x)(b\circ a) + b\circ(r_{\mathcal{A}}(x)a) + l_{\mathcal{B}}(b)(l_{\mathcal{B}}(a)x) + r_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b\}$

(3.13) $\Leftrightarrow r_{\mathcal{A}}(x)([a,b]) = r_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a + r_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b + a\circ(r_{\mathcal{A}}(x)b) + b\circ(r_{\mathcal{A}}(x)a)$ since (3.2) hold,

(3.13) $\Leftrightarrow$ (3.5) and $l_{\mathcal{B}}$ is a linear representation of the sub-adjacent Mock-Lie algebra $\mathcal{G}(\mathcal{B})$.

Hence, $\mathcal{A} \bowtie^{-1} \mathcal{B}$ is a quasi-antiassociative algebra if and only if $(l_{\mathcal{A}}, r_{\mathcal{A}})$ is a bimodule of $\mathcal{A}$ and $(l_{\mathcal{B}}, r_{\mathcal{B}})$ is a bimodule of $\mathcal{B}$ and equations (3.5) – (3.8) hold.

On the other hand, if $\mathcal{A}$ and $\mathcal{B}$ are quasi-antiassociative sub-algebras of a quasi-antiassociative algebra C such that $C = \mathcal{A} \oplus \mathcal{B}$ which is a direct sum of the underlying vector spaces of $\mathcal{A}$ and $\mathcal{B}$, then the linear maps

$$l_{\mathcal{A}}, r_{\mathcal{A}} : \mathcal{A} \to \mathfrak{gl}(\mathcal{B}), \quad l_{\mathcal{B}}, r_{\mathcal{B}} : \mathcal{B} \to \mathfrak{gl}(\mathcal{A}).$$

defined by

$$x * a = l_{\mathcal{A}}(x)a + r_{\mathcal{B}}(a)x$$
$$a * x = l_{\mathcal{B}}(a)x + r_{\mathcal{A}}(x)a$$

satisfy the equations (3.5) – (3.8). In addition, $(l_{\mathcal{A}}, r_{\mathcal{A}})$ is a bimodule of $\mathcal{A}$ and $(l_{\mathcal{B}}, r_{\mathcal{B}})$ is a bimodule of $\mathcal{B}$. □

Corollary 3.7. *Let $(\mathcal{A}, \mathcal{B}, l_{\mathcal{A}}, r_{\mathcal{A}}, l_{\mathcal{B}}, r_{\mathcal{B}})$ be a matched pair of quasi-antiassociative algebras. Then, $(\mathcal{G}(\mathcal{A}), \mathcal{G}(\mathcal{B}), l_{\mathcal{A}} + r_{\mathcal{A}}, l_{\mathcal{B}} + r_{\mathcal{B}})$ is a matched pair of sub-adjacent Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{B})$.*

Proof. By using the Proposition 3.3 and the bimodules $(l_{\mathcal{A}}, r_{\mathcal{A}}, \mathcal{B})$ and $(l_{\mathcal{B}}, r_{\mathcal{B}}, \mathcal{A})$, we have: $\mathrm{ad}_{\mathcal{A}} := l_{\mathcal{A}} + r_{\mathcal{A}}$ and $\mathrm{ad}_{\mathcal{B}} := l_{\mathcal{B}} + r_{\mathcal{B}}$ are the linear representations of the sub-adjacent Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{B})$ of the quasi-antiassociative algebras $\mathcal{A}$ and $\mathcal{B}$, respectively. Then, the statement that $\mathcal{G}(\mathcal{A}) \bowtie^{-1,\mathrm{ad}_{\mathcal{A}}}_{\mathrm{ad}_{\mathcal{B}}} \mathcal{G}(\mathcal{B})$ is a matched pair of the Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{B})$ follows from Theorem 3.6. By analogous step giving:

$$\mathrm{ad}_{\mathcal{A}}(x)\,[a,b] - [\mathrm{ad}_{\mathcal{A}}(x)a, b] - [a, \mathrm{ad}_{\mathcal{A}}(x)b] - \mathrm{ad}_{\mathcal{A}}(\mathrm{ad}_{\mathcal{B}}(a)x)b - \mathrm{ad}_{\mathcal{A}}(\mathrm{ad}_{\mathcal{B}}(b)x)a = 0, \tag{3.14}$$

$$\mathrm{ad}_{\mathcal{B}}(a)\,[x,y] - [\mathrm{ad}_{\mathcal{B}}(a)x, y] - [x, \mathrm{ad}_{\mathcal{B}}(a)y]$$

$$-\mathrm{ad}_{\mathcal{B}}(\mathrm{ad}_{\mathcal{A}}(x)a)y - \mathrm{ad}_{\mathcal{B}}(\mathrm{ad}_{\mathcal{A}}(y)a)x = 0. \tag{3.15}$$

Thus, we first have:

$$\begin{aligned}
&\mathrm{ad}_{\mathcal{A}}(x)[a,b]-[\mathrm{ad}_{\mathcal{A}}(x)a,b]-[a,\mathrm{ad}_{\mathcal{A}}(x)b]-\mathrm{ad}_{\mathcal{A}}(\mathrm{ad}_{\mathcal{B}}(a)x)b-\mathrm{ad}_{\mathcal{A}}(\mathrm{ad}_{\mathcal{B}}(b)x)a\\
&= (l_{\mathcal{A}}+r_{\mathcal{A}})(x)\,[a,b] - [(l_{\mathcal{A}}+r_{\mathcal{A}})(x)a,b] - [a,(l_{\mathcal{A}}+r_{\mathcal{A}})(x)b]\\
&\quad -(l_{\mathcal{A}}+r_{\mathcal{A}})((l_{\mathcal{B}}+r_{\mathcal{B}})(a)x)b - (l_{\mathcal{A}}+r_{\mathcal{A}})((l_{\mathcal{B}}+r_{\mathcal{B}})(b)x)a\\
&= l_{\mathcal{A}}(x)[a,b] + r_{\mathcal{A}}(x)[a,b] - [l_{\mathcal{A}}(x)a,b] - [r_{\mathcal{A}}(x)a,b] - [a,l_{\mathcal{A}}(x)b]\\
&\quad - [a,r_{\mathcal{A}}(x)b] - l_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b - l_{\mathcal{A}}(r_{\mathcal{B}}(a)x)b - r_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b - r_{\mathcal{A}}(r_{\mathcal{B}}(a)x)b\\
&\quad -l_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a - l_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a - r_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a - r_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a\\
&= l_{\mathcal{A}}(x)(a\circ b) + l_{\mathcal{A}}(x)(b\circ a) + r_{\mathcal{A}}(x)(a\circ b) + r_{\mathcal{A}}(x)(b\circ a) - (l_{\mathcal{A}}(x)a)\circ b\\
&\quad -b\circ(l_{\mathcal{A}}(x)a) - (r_{\mathcal{A}}(x)a)\circ b - b\circ(r_{\mathcal{A}}(x)a) - a\circ(l_{\mathcal{A}}(x)b) - (l_{\mathcal{A}}(x)b)\circ a\\
&\quad -a\circ(r_{\mathcal{A}}(x)b) - (r_{\mathcal{A}}(x)b)\circ a - l_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b - l_{\mathcal{A}}(r_{\mathcal{B}}(a)x)b - r_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b\\
&\quad -r_{\mathcal{A}}(r_{\mathcal{B}}(a)x)b - l_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a - l_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a - r_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a - r_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a\\
&= \{r_{\mathcal{A}}(x)([a,b]) - r_{\mathcal{A}}(l_{\mathcal{B}}(b)x)a - r_{\mathcal{A}}(l_{\mathcal{B}}(a)x)b - a\circ(r_{\mathcal{A}}(x)b) - b\circ(r_{\mathcal{A}}(x)a)\}\\
&\quad +\{-l_{\mathcal{A}}(x)(a\circ b) - l_{\mathcal{A}}(l_{\mathcal{B}}(a)x - r_{\mathcal{B}}(a)x)b - (l_{\mathcal{A}}(x)a - r_{\mathcal{A}}(x)a)\circ b\\
&\quad -r_{\mathcal{A}}(r_{\mathcal{B}}(b)x)a - a\circ(l_{\mathcal{A}}(x)b)\}\\
&\quad +\{-l_{\mathcal{A}}(x)(b\circ a) - l_{\mathcal{A}}(l_{\mathcal{B}}(b)x - r_{\mathcal{B}}(b)x)a - (l_{\mathcal{A}}(x)b - r_{\mathcal{A}}(x)b)\circ a\\
&\quad -r_{\mathcal{A}}(r_{\mathcal{B}}(a)x)b - b\circ(l_{\mathcal{A}}(x)a)\} = 0.
\end{aligned}$$

Secondly:

$$\begin{aligned}
&\mathrm{ad}_{\mathcal{B}}(a)[x,y]-[\mathrm{ad}_{\mathcal{B}}(a)x,y]-[x,\mathrm{ad}_{\mathcal{B}}(a)y]-\mathrm{ad}_{\mathcal{B}}(\mathrm{ad}_{\mathcal{A}}(x)a)y-\mathrm{ad}_{\mathcal{B}}(\mathrm{ad}_{\mathcal{A}}(y)a)x\\
&= (l_{\mathcal{B}}+r_{\mathcal{B}})(a)\,[x,y] - [(l_{\mathcal{B}}+r_{\mathcal{B}})(a)x,y] - [x,(l_{\mathcal{B}}+r_{\mathcal{B}})(a)y]\\
&\quad -(l_{\mathcal{B}}+r_{\mathcal{B}})((l_{\mathcal{A}}+r_{\mathcal{A}})(x)a)y - (l_{\mathcal{B}}+r_{\mathcal{B}})((l_{\mathcal{A}}+r_{\mathcal{A}})(y)a)x\\
&= l_{\mathcal{B}}(a)[x,y] + r_{\mathcal{B}}(a)[x,y] - [l_{\mathcal{B}}(a)x,y] - [r_{\mathcal{B}}(a)x,y] - [x,l_{\mathcal{B}}(a)y]\\
&\quad - [x,r_{\mathcal{B}}(a)y] - l_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y - l_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y - r_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y - r_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y\\
&\quad -l_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x - l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x - r_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x - r_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x\\
&= \{r_{\mathcal{A}}(a)[x,y] - r_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x - r_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y - x(r_{\mathcal{B}}(a)y) - y(r_{\mathcal{B}}(a)x)\}\\
&\quad +\{-l_{\mathcal{B}}(a)(xy) - l_{\mathcal{B}}(l_{\mathcal{A}}(x)a)y - (r_{\mathcal{B}}(a)x)y - x(l_{\mathcal{B}}(a)y) - r_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x\\
&\quad -(l_{\mathcal{B}}(a)x)y - l_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y - l_{\mathcal{B}}(a)(xy)\} + \{-l_{\mathcal{B}}(a)(yx) - l_{\mathcal{B}}(l_{\mathcal{A}}(y)a)x\\
&\quad -(r_{\mathcal{B}}(a)y)x - y(l_{\mathcal{B}}(a)x) - r_{\mathcal{B}}(r_{\mathcal{A}}(x)a)y - (l_{\mathcal{B}}(a)y)x - l_{\mathcal{B}}(r_{\mathcal{A}}(y)a)x\\
&\quad -l_{\mathcal{B}}(a)(yx)\} = 0.
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
&\mathrm{ad}_{\mathcal{A}}(x)\,[a,b] - [\mathrm{ad}_{\mathcal{A}}(x)a,b] - [a,\mathrm{ad}_{\mathcal{A}}(x)b]\\
&\qquad\qquad - \mathrm{ad}_{\mathcal{A}}(\mathrm{ad}_{\mathcal{B}}(a)x)b - \mathrm{ad}_{\mathcal{A}}(\mathrm{ad}_{\mathcal{B}}(b)x)a = 0,\\
&\mathrm{ad}_{\mathcal{B}}(a)\,[x,y] - [\mathrm{ad}_{\mathcal{B}}(a)x,y] - [x,\mathrm{ad}_{\mathcal{B}}(a)y]\\
&\qquad\qquad - \mathrm{ad}_{\mathcal{B}}(\mathrm{ad}_{\mathcal{A}}(x)a)y - \mathrm{ad}_{\mathcal{B}}(\mathrm{ad}_{\mathcal{A}}(y)a)x = 0.
\end{aligned}$$

Hence, $(\mathcal{G}(\mathcal{A}),\mathcal{G}(\mathcal{B}),\mathrm{ad}_{\mathcal{A}},\mathrm{ad}_{\mathcal{B}})$ is a matched pair of sub-adjacent Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{B})$. □

Definition 3.8. Let (l, r, V) be a bimodule of a quasi-antiassociative algebra $\mathcal{A}$, where V is a finite dimensional vector space. The dual maps l^*, r^* of the linear maps l, r, are defined, respectively, as: $l^*, r^* : \mathcal{A} \to \mathfrak{gl}(V^*)$ such that: for all $x \in \mathcal{A}, u^* \in V^*, v \in V$,

$$\begin{array}{ccccccccc} l^* : \mathcal{A} & \longrightarrow & \mathfrak{gl}(V^*) & & & & & & \\ & & & V^* & \longrightarrow & V^* & & & \\ x & \longmapsto & l_x^* : & u^* & \longmapsto & l_x^* u^* : & V & \longrightarrow & \mathbb{K} \\ & & & & & & v & \longmapsto & \langle l_x^* u^*, v\rangle := \langle u^*, l_x v\rangle , \end{array} \tag{3.16}$$

$$\begin{array}{ccccccccc} r^* : \mathcal{A} & \longrightarrow & \mathfrak{gl}(V^*) & & & & & & \\ & & & V^* & \longrightarrow & V^* & & & \\ x & \longmapsto & r_x^* : & u^* & \longmapsto & r_x^* u^* : & V & \longrightarrow & \mathbb{K} \\ & & & & & & v & \longmapsto & \langle r_x^* u^*, v\rangle := \langle u^*, -r_x v\rangle . \end{array} \tag{3.17}$$

Proposition 3.9. *Let $(\mathcal{A}, \cdot)$ be a quasi-antiassociative algebra and l, r be two linear maps $\mathcal{A} \to \mathfrak{gl}(V)$, where V is a finite dimensional vector space. The following conditions are equivalent:*

1. *(l, r, V) is a bimodule of $\mathcal{A}$.*
2. *$(l^* + r^*, -r^*, V^*)$ is a bimodule of $\mathcal{A}$.*

Proof. (1) $\Rightarrow$ (2): Suppose that (l, r, V) is a bimodule of $(\mathcal{A}, \cdot)$ and show that $(l^* + r^*, -r^*, V^*)$ is also a bimodule of $(\mathcal{A}, \cdot)$. We have: firstly,

$$\begin{aligned} \langle [(l^*+r^*)_{xy} + (l^*+r^*)_x(l^*+r^*)_y]u^*, v\rangle &= \langle (l^*+r^*)_{xy}u^*, v\rangle + \langle [(l^*+r^*)_x(l^*+r^*)_y]u^*, v\rangle \\ &= -\langle (l+r)_{xy}(v), u^*\rangle - \langle [(l+r)_y(l+r)_x](v), u^*\rangle \\ &= -\langle (l^*+r^*)_{yx}u^*, v\rangle - \langle (l^*+r^*)_y(l^*+r^*)_x)u^*, v\rangle. \end{aligned}$$

Therefore,

$$(l^*+r^*)_{xy} + (l^*+r^*)_x(l^*+r^*)_y = -(l^*+r^*)_{yx} - (l^*+r^*)_y(l^*+r^*)_x, \ \forall\, x, y \in \mathcal{A} \tag{3.18}$$

Secondly,

$$\begin{aligned} \langle r_y^*(l^*+r^*)_x u^* + (l^*+r^*)_x r_y^* u^*, v\rangle &= \langle (l+r)_x r_y v, u^*\rangle + \langle r_y(l+r)_x v, u^*\rangle \\ &= \langle l_x(r_y(v)) + r_x(r_y(v)), u^*\rangle + \langle r_y(l_x(v)) + r_y(r_x(v)), u^*\rangle \\ &= \langle r_x(r_y(v)) + r_{xy}v, u^*\rangle \\ &= \big\langle (r_y^* r_x^* + r_{xy}^*)u^*, v\big\rangle. \end{aligned}$$

Therefore

$$(-r^*)_y(-r^*)_x + (-r^*)_{xy} = -[(l^* + r^*)_y, -r_x^*], \ \forall\, x, y \ \in \mathcal{A}. \tag{3.19}$$

By considering the relations (3.18) and (3.19), we conclude that $(l^* + r^*, -r^*, V^*)$ is a bimodule of $(\mathcal{A}, \cdot)$.

(2) $\Rightarrow$ (1): The converse, (i.e., by supposing that $(l^* + r^*, -r^*, V^*)$ is a bimodule of $(\mathcal{A}, \cdot)$ then (l, r, V) is also a bimodule of $(\mathcal{A}, \cdot)$), can be proved by direct calculations by using similar relations as for the first part of the proof. □

Example 3.10. Let $\mathcal{A}$ be a quasi-antiassociative algebra. Then $(L, 0)$, $(\mathrm{ad}, 0)$ and (L, R) are bimodules of $\mathcal{A}$, where $\mathrm{ad} = L + R : \mathcal{A} \to gl(\mathcal{A})$ is the adjoint representation of the Mock Lie algebra $\mathcal{G}(\mathcal{A})$. On the other hand, $(L^*, 0)$, $(\mathrm{ad}^*, 0)$ and $(\mathrm{ad}^*, -R^*)$ are bimodules of $\mathcal{A}$, too, where ad^* and L^* are the dual representations of the adjoint representation and the regular representation of the sub-adjacent Mock Lie algebra $\mathcal{G}(\mathcal{A})$ respectively.

Theorem 3.11. *Let $(\mathcal{A}, \cdot)$ be a quasi-antiassociative algebra. Suppose that there exists a quasi-antiassociative algebra structure " $\circ$ " on its dual space $\mathcal{A}^*$. Then, $(\mathcal{A}, \mathcal{A}^*, \mathrm{ad}^*_\cdot, -R^*_\cdot, \mathrm{ad}^*_\circ, -R^*_\circ)$ is a matched pair of quasi-antiassociative algebras $\mathcal{A}$ and $\mathcal{A}^*$ if and only if $(\mathcal{G}(\mathcal{A}), \mathcal{G}(\mathcal{A}^*), L^*_\cdot, L^*_\circ)$ is a matched pair of Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{A}^*)$.*

Proof. By considering the Corollary 3.7, setting $l_{\mathcal{A}} := \mathrm{ad}^*_\cdot = L^*_\cdot + R^*_\cdot, r_{\mathcal{A}} := -R^*_\cdot, l_{\mathcal{B}} := \mathrm{ad}^*_\circ = L^*_\circ + R^*_\circ, r_{\mathcal{B}} := -R^*_\circ$, and exploiting the Definition 3.5 with $\mathcal{G} := \mathcal{G}(\mathcal{A}), \mathcal{H} := \mathcal{G}(\mathcal{A}^*), \rho := L^*_\cdot, \mu := L^*_\circ$, the proof is established. □

4. Quasi-antiassociative bialgebras

Definition 4.1. [2] Let $(\mathcal{G}, [,])$ be a Mock-Lie algebra. A 1-cocycle δ is associated to a representation (ρ, V) of $(\mathcal{G}, [,])$, if the equation

$$\delta([x, y]) = -\rho(x)\delta(y) - \rho(y)\delta(x), \quad \forall x, y \in \mathcal{G}, \tag{4.1}$$

is satisfied.

Proposition 4.2. *Let $(\mathcal{G}, [,])$ be a Mock-Lie algebra. Suppose $\rho : \mathcal{G} \to \mathfrak{gl}(V)$ and $\mu : \mathcal{G} \to \mathfrak{gl}(W)$ be two linear representations of $\mathcal{G}$, where V and W are two vector spaces. Then, the linear map $\rho \otimes 1 + 1 \otimes \mu : \mathcal{G} \to \mathfrak{gl}(V \otimes W)$ given by $(\rho \otimes 1 + 1 \otimes \mu)(x)(v, w) := \rho(x)v \otimes w + v \otimes \mu(x)w$ is also a representation of $\mathcal{G}$.*

Proof. Let $(\mathcal{G}, [,])$ be a Mock-Lie algebra, $(\mathcal{G}, \rho)$ and $(\mathcal{G}, \mu)$ be two representations of $(\mathcal{G}, [,])$. For all $x, y \in \mathcal{G}$ and $v \otimes w \in V \otimes W$, we have

$$(\rho \otimes 1 + 1 \otimes \mu)([x, y]) = \rho([x, y]) \otimes 1 + 1 \otimes \mu([x, y])$$

$$
\begin{aligned}
&= -\rho(x)\rho(y)\otimes 1 - 1\otimes\mu(x)\mu(y)\\
&\quad -1\otimes\mu(y)\mu(x) - \rho(y)\rho(x)\otimes 1\\
&= -\rho(x)\rho(y)v\otimes 1 - 1\otimes\mu(x)\mu(y)\\
&\quad -1\otimes\mu(y)\mu(x) - \rho(y)\rho(x)\otimes 1\\
&\quad -\rho(x)\otimes\mu(y) - \rho(x)\otimes\mu(y)\\
&\quad -\rho(y)\otimes\mu(x) - \rho(y)\otimes\mu(x)\\
&= -[\rho(x)\otimes 1, \rho(y)\otimes 1] - [\rho(x)\otimes 1, 1\otimes\mu(y)]\\
&\quad -[1\otimes\mu(x), \rho(y)\otimes 1] - [1\otimes\mu(x), \rho(y)\otimes 1]\\
&= -[\rho(x)\otimes 1 + 1\otimes\mu(x), \rho(y)\otimes 1 + 1\otimes\mu(y)].
\end{aligned}
$$

Therefore, we have $\forall x, y \in \mathcal{G}$;

$$(\rho\otimes 1 + 1\otimes\mu)([x,y]) = -[\rho(x)\otimes 1 + 1\otimes\mu(x), \rho(y)\otimes 1 + 1\otimes\mu(y)],$$

which exactly means that $\rho\otimes 1 + 1\otimes\mu$ is a representation of $(\mathcal{G};[,])$ on $V\otimes W$. □

Theorem 4.3. *Let $\mathcal{A}$ be a quasi-antiassociative algebra with the product given by the linear map $\beta^* : \mathcal{A}\otimes\mathcal{A}\to\mathcal{A}$. Suppose there is a quasi-antiassociative algebra structure " $\circ$ " on the dual space $\mathcal{A}^*$ provided by a linear map $\alpha^* : \mathcal{A}^*\otimes\mathcal{A}^*\to\mathcal{A}^*$. Then, $(\mathcal{G}(\mathcal{A}), \mathcal{G}(\mathcal{A}^*), L^*_\cdot, L^*_\circ)$ is a matched pair of Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{A}^*)$ if and only if $\alpha : \mathcal{A}\to\mathcal{A}\otimes\mathcal{A}$ is an antiderivation of $\mathcal{G}(\mathcal{A})$ associated to $.\otimes 1 + 1\otimes\mathrm{ad}.$ and $\beta : \mathcal{A}^*\to\mathcal{A}^*\otimes\mathcal{A}^*$ is an antiderivation of $\mathcal{G}(\mathcal{A}^*)$ associated to $L_\circ\otimes 1 + 1\otimes\mathrm{ad}_\circ$.*

Proof. Let $\{e_1, e_2, \cdots, e_n\}$ be a basis of $\mathcal{A}$ and $\{e_1^*, e_2^*, \cdots, e_n^*\}$ its dual basis. Consider $e_i\cdot e_j = \sum_{k=1}^n c_{ij}^k e_k$ and $e_i^*\circ e_j^* = \sum_{k=1}^n f_{ij}^k e_k^*$, where $c_{ij}^k, f_{ij}^k \in \mathbb{K}$ are structure constants associated to $\cdot$ and $\circ$, respectively. Then, it follows that:

$$\alpha(e_k) = \sum_{i,j=1}^n f_{ij}^k e_i\otimes e_j, \quad \beta(e_k^*) = \sum_{i,j=1}^n c_{ij}^k e_i^*\otimes e_j^*, \quad \text{and}$$

$$\alpha([e_i,e_j]) = \sum_{m,l=1}^n\sum_{k=1}^n \left\{(c_{ij}^k + c_{ji}^k) f_{ml}^k\right\} e_m\otimes e_l, \tag{4.2}$$

$$\beta([e_i^*,e_j^*]) = \sum_{m,l=1}^n\sum_{k=1}^n \left\{(f_{ij}^k + f_{ji}^k) c_{ml}^k\right\} e_m^*\otimes e_l^*. \tag{4.3}$$

Now, let us compute the quantities:

$$\{(-L.)(e_i)\otimes 1 + 1\otimes(-\mathrm{ad}.)(e_i)\}\alpha(e_j) + \{(-L.)(e_j)\otimes 1 + 1\otimes(-\mathrm{ad}.)(e_j)\}\alpha(e_i)$$

and

$$\{(-L_\circ)(e_i^*)\otimes 1+1\otimes(-\operatorname{ad}_\circ)(e_i^*)\}\beta(e_j^*)+\{(-L_\circ)(e_j^*)\otimes 1+1\otimes(-\operatorname{ad}_\circ)(e_j^*)\}\beta(e_i^*).$$

We have:

$$\begin{aligned}
&\{(-L.)(e_i)\otimes 1+1\otimes(-\mathrm{ad}.)(e_i)\}\alpha(e_j)+\{(-L.)(e_j)\otimes 1+1\otimes(-\mathrm{ad}.)(e_j)\}\alpha(e_i)\\
&=\{-(L.(e_i)\otimes 1)\alpha(e_j)-(L.(e_j)\otimes 1)\alpha(e_i)\}+\{-(1\otimes\mathrm{ad}.(e_j))\alpha(e_i)\\
&-(1\otimes\mathrm{ad}.(e_i))\alpha(e_j)\}=\sum_{k,l=1}^{n}\{-f_{kl}^j(L.(e_i)\otimes 1)(e_k\otimes e_l)-f_{kl}^i(L.(e_j)\otimes 1)(e_k\otimes e_l)\}\\
&+\sum_{m,k=1}^{n}\{-f_{mk}^j(1\otimes\mathrm{ad}.(e_i))(e_m\otimes e_k)-f_{mk}^i(1\otimes\mathrm{ad}.(e_j))(e_m\otimes e_k)\}\\
&=\sum_{k,l=1}^{n}\{-f_{kl}^j([e_i,e_k]\otimes e_l)-f_{kl}^i([e_j,e_k]\otimes e_l)\}\\
&+\sum_{m,k=1}^{n}\{-f_{mk}^j(e_m\otimes[e_i,e_k])-f_{mk}^i(e_m\otimes[e_j,e_k])\}\\
&=\sum_{k,l=1}^{n}\left\{-f_{kl}^j\left(\sum_{m=1}^{n}(c_{ik}^m+c_{ki}^m)(e_m\otimes e_l)\right)-f_{kl}^i\left(\sum_{m=1}^{n}(c_{jk}^m+c_{kj}^m)(e_m\otimes e_l)\right)\right\}\\
&+\sum_{m,k=1}^{n}\left\{-f_{mk}^j\left(\sum_{l=1}^{n}(c_{ik}^l+c_{ki}^l)(e_m\otimes e_l)\right)-f_{mk}^i\left(\sum_{l=1}^{n}(c_{jk}^l+c_{kj}^l)(e_m\otimes e_l)\right)\right\}\\
&=\sum_{m,l=1}^{n}\sum_{k=1}^{n}\Big\{-f_{kl}^j(c_{ik}^m+c_{ki}^m)-f_{kl}^i(c_{jk}^m+c_{kj}^m)-f_{mk}^j(c_{ik}^l+c_{ki}^l)-f_{mk}^i(c_{jk}^l+c_{kj}^l)\Big\}(e_m\otimes e_l).
\end{aligned}$$

Therefore, we have:

$$\begin{aligned}
&\{(-L.)(e_i)\otimes 1+1\otimes(-\operatorname{ad}.)(e_i)\}\alpha(e_j)+\{(-L.)(e_j)\otimes 1+1\otimes(-\operatorname{ad}.)(e_j)\}\alpha(e_i)\\
&=\sum_{m,l=1}^{n}\sum_{k=1}^{n}\{-f_{kl}^j(c_{ik}^m+c_{ki}^m)-f_{kl}^i(c_{jk}^m+c_{kj}^m)-f_{mk}^j(c_{ik}^l+c_{ki}^l)\\
&-f_{mk}^i(c_{jk}^l+c_{kj}^l)\}e_m\otimes e_l.
\end{aligned}\tag{4.4}$$

Taking into account the fact that α is an antiderivation of $\mathcal{G}(\mathcal{A})$ associated to $L.\otimes 1+1\otimes\mathrm{ad}.$, and using the relations (4.2) and (4.4) yield:

$$\sum_{k=1}^{n}(c_{ij}^k+c_{ji}^k)f_{ml}^k=\sum_{k=1}^{n}\{-f_{kl}^i(c_{jk}^m+c_{kj}^m)-f_{kl}^j(c_{ik}^m+c_{ki}^m)-f_{mk}^i(c_{jk}^l+c_{kj}^l)-f_{mk}^j(c_{ik}^l+c_{ki}^l)\}.\tag{4.5}$$

Besides, we obtain:

$$\{(-L_\circ)(e_i^*)\otimes 1+1\otimes(-\operatorname{ad}_\circ)(e_i^*)\}\beta(e_j^*)+\{(-L_\circ)(e_j^*)\otimes 1+1\otimes(-\operatorname{ad}_\circ)(e_j^*)\}\beta(e_i^*)$$

$$
\begin{aligned}
&= \{(-L_\circ)(e_i^*)\otimes 1)\beta(e_j^*) + (-L_\circ)(e_j^*\otimes 1)\beta(e_i^*)\}\\
&\quad +\{(1\otimes(-\operatorname{ad}_\circ)(e_i^*))\beta(e_j^*) + (1\otimes(-\operatorname{ad}_\circ)(e_j^*))\beta(e_i^*)\}\\
&= \sum_{k,l=1}^{n}\{-c_{kl}^j(L_\circ(e_i^*)\otimes 1) - c_{kl}^i(L_\circ(e_j^*)\otimes 1)\}(e_k^*\otimes e_l^*)\\
&\quad + \sum_{m,k=1}^{n}\{-c_{mk}^j(1\otimes\operatorname{ad}_\circ(e_i^*)) - c_{mk}^i(1\otimes\operatorname{ad}_\circ(e_j^*))\}(e_m^*\otimes e_k^*)\\
&= \sum_{k,l=1}^{n}\Big\{-c_{kl}^j[e_i^*,e_k^*] - c_{kl}^i[e_j^*,e_k^*]\Big\}\otimes e_l^* + \sum_{m,k=1}^{n} e_m^*\otimes\Big\{-c_{mk}^j[e_i^*,e_k^*] - c_{mk}^i[e_j^*,e_k^*]\Big\}\\
&= \sum_{k,l=1}^{n}\sum_{m=1}^{n}\Big\{-c_{kl}^j(f_{ik}^m + f_{ki}^m) - c_{kl}^i(f_{jk}^m + f_{kj}^m)\Big\}(e_m^*\otimes e_l^*)\\
&\quad + \sum_{m,k=1}^{n}\sum_{l=1}^{n}\Big\{-c_{mk}^j(f_{ik}^l + f_{ki}^l) - c_{mk}^i(f_{jk}^l + f_{kj}^l)\Big\}(e_m^*\otimes e_l^*)\\
&= \sum_{m,l=1}^{n}\sum_{k=1}^{n}\Big\{-c_{kl}^j(f_{ik}^m+f_{ki}^m) - c_{kl}^i(f_{jk}^m+f_{kj}^m) - c_{mk}^j(f_{ik}^l+f_{ki}^l) - c_{mk}^i(f_{jk}^l+f_{kj}^l)\Big\}(e_m^*\otimes e_l^*).
\end{aligned}
$$

Thus

$$
\begin{aligned}
&\{(-L_\circ)(e_i^*)\otimes 1+1\otimes(-\operatorname{ad}_\circ)(e_i^*)\}\beta(e_j^*)+\{(-L_\circ)(e_j^*)\otimes 1+1\otimes(-\operatorname{ad}_\circ)(e_j^*)\}\beta(e_i^*)\\
&=\sum_{m,l=1}^{n}\sum_{k=1}^{n}\{-c_{kl}^j(f_{ik}^m+f_{ki}^m)-c_{kl}^i(f_{jk}^m+f_{kj}^m)-c_{mk}^j(f_{ik}^l+f_{ki}^l)-c_{mk}^i(f_{jk}^l+f_{kj}^l)\}(e_m^*\otimes e_l^*). \quad (4.6)
\end{aligned}
$$

Since β is the antiderivation issued from $L_\circ\otimes 1 + 1\otimes\operatorname{ad}_\circ$, using the relations (4.3) and (4.6) we obtain:

$$
\sum_{k=1}^{n}(f_{ij}^k+f_{ji}^k)c_{ml}^k = \sum_{k=1}^{n}\{-c_{kl}^i(f_{jk}^m+f_{kj}^m)-c_{kl}^j(f_{ik}^m+f_{ki}^m)-c_{mk}^i(f_{jk}^l+f_{kj}^l)-c_{mk}^j(f_{ik}^l+f_{ki}^l)\}. \quad (4.7)
$$

Now, let us find the relations associated to the equations (3.3)-(3.4) of the matched pair of Mock-Lie algebras $\mathcal{G}(\mathcal{A})$ and $\mathcal{G}(\mathcal{A}^*)$. We have $\forall\, i,j,k$:

$$
\langle(-L_\cdot^*)(e_i)e_j^*, e_k\rangle = -\langle e_j^*, L_\cdot(e_i)e_k\rangle = -\langle e_j^*,[e_i,e_k]\rangle = -\left\langle e_j^*, \sum_{m=1}^{n}(c_{ik}^m+c_{ki}^m)e_m\right\rangle = -(c_{ik}^j+c_{ki}^j),
$$

$$
\langle(-L_\cdot^*)(e_i)e_j^*, e_k\rangle = -\left\langle\sum_{k=1}^{m}(c_{ik}^j+c_{ki}^j)e_j^*, e_k\right\rangle,
$$

providing

$$
(-L_\cdot^*)(e_i)e_j^* = -\sum_{k=1}^{n}(c_{ik}^j+c_{ki}^j)e_k^*.
$$

Similarly,

$$
(-L_\circ^*)(e_i^*)e_j = -\sum_{k=1}^{n}(f_{ik}^j+f_{ki}^j)e_k,
$$

$$
(-L_\circ^*)(e_m^*)[e_i,e_j] = \sum_{k=1}^{n}(c_{ij}^k - c_{ji}^k)(-L_\circ^*)(e_m^*)e_k - \sum_{l=1}^{n}\sum_{k=1}^{n}(c_{ij}^k - c_{ji}^k)(f_{ml}^k - f_{lm}^k)e_l,
$$

and,

$$(-L_\circ^*)(e_m^*)[e_i,e_j] = -\sum_{l=1}^{n}\sum_{k=1}^{n}(c_{ij}^k+c_{ji}^k)(f_{ml}^k+f_{lm}^k)e_l.$$

Then

$$\begin{aligned}
&- L_\circ^*(L_\cdot^*(e_i)e_m^*)e_j+[e_i,L_\circ^*(e_m^*)e_j]-L_\circ^*(L_\cdot^*(e_j)e_m^*)e_i+[L_\circ^*(e_m^*)e_i,e_j]\\
&-\sum_{k=1}^{n}(c_{ik}^m+c_{ki}^m)L_\circ^*(e_k^*)e_j-\sum_{k=1}^{n}(f_{mk}^j+f_{km}^j)[e_i,e_k]\\
&-\sum_{k=1}^{n}(c_{jk}^m+c_{kj}^m)L_\circ^*(e_k^*)e_i-\sum_{k=1}^{n}(f_{mk}^i+f_{km}^i)[e_k,e_j]\\
&=-\sum_{k=1}^{n}\sum_{l=1}^{n}\Big\{(c_{ik}^m+c_{ki}^m)(f_{kl}^j+f_{lk}^j)e_l+(f_{mk}^j+f_{km}^j)(c_{ik}^l+c_{ki}^l)e_l\Big\}\\
&-\sum_{k=1}^{n}\sum_{l=1}^{n}\Big\{(c_{jk}^m+c_{kj}^m)(f_{kl}^i+f_{lk}^i)e_l+(f_{mk}^i+f_{km}^i)(c_{kj}^l+c_{jk}^l)e_l\Big\}\\
&=-\sum_{l=1}^{n}\sum_{k=1}^{n}\{(c_{ik}^m+c_{ki}^m)(f_{kl}^j+f_{lk}^j)+(f_{mk}^j+f_{km}^j)(c_{ik}^l+c_{ki}^l)\\
&+(c_{jk}^m+c_{kj}^m)(f_{kl}^i+f_{lk}^i)+(f_{mk}^i+f_{km}^i)(c_{kj}^l+c_{jk}^l)\}e_l.
\end{aligned}$$

Using the fact that $(\mathcal{G}(\mathcal{A}),\mathcal{G}(\mathcal{A}^*),L_\cdot^*,L_\circ^*)$ is a matched pair of Mock-Lie algebras, we have

$$\begin{aligned}
\sum_{k=1}^{n}(c_{ij}^k+c_{ji}^k)(f_{ml}^k+f_{lm}^k) = &-\sum_{k=1}^{n}\{(c_{ik}^m+c_{ki}^m)(f_{kl}^j+f_{lk}^j)+(f_{mk}^j+f_{km}^j)(c_{ik}^l+c_{ki}^l)\\
&+(c_{jk}^m+c_{kj}^m)(f_{kl}^i+f_{lk}^i)+(f_{mk}^i+f_{km}^i)(c_{jk}^l+c_{kj}^l)\},
\end{aligned}$$

that is,

$$\begin{aligned}
&\sum_{k=1}^{n}(c_{ij}^k+c_{ji}^k)f_{ml}^k+\sum_{k=1}^{n}(c_{ik}^m+c_{ki}^m)f_{kl}^j+(c_{ik}^l+c_{ki}^l)f_{mk}^j+(c_{jk}^m+c_{kj}^m)f_{kl}^i+(c_{jk}^l+c_{kj}^l)f_{mk}^i\\
&=-\sum_{k=1}^{n}(c_{ij}^k+c_{ji}^k)f_{lm}^k-\sum_{k=1}^{n}(c_{ik}^m+c_{ki}^m)f_{lk}^j-(c_{ik}^l+c_{ki}^l)f_{km}^j-(c_{jk}^m+c_{kj}^m)f_{lk}^i-(c_{jk}^l+c_{kj}^l)f_{km}^i.
\end{aligned}$$

Replacing l (resp. m) by m (resp. l) in the right-hand side of the equality leads to:

$$\sum_{k=1}^{n}(c_{ij}^k+c_{ji}^k)f_{ml}^k=\sum_{k=1}^{n}\{-(c_{ik}^m+c_{ki}^m)f_{kl}^j-(c_{ik}^l+c_{ki}^l)f_{mk}^j-(c_{jk}^m+c_{kj}^m)f_{kl}^i-(c_{jk}^l+c_{kj}^l)f_{mk}^i\},$$

which is identical to the equation (4.5). Besides,

$$(-\operatorname{ad}_\cdot^*)(e_m)[e_i^*,e_j^*] = -\sum_{l=1}^{n}\sum_{k=1}^{n}\Big\{(f_{ij}^k+f_{ji}^k)(c_{ml}^k+c_{lm}^k)\Big\}e_l^*$$

$$\begin{aligned}
&-\operatorname{ad}_\cdot^*(\operatorname{ad}_\circ^*(e_i^*)e_m)e_j^*+[e_i^*,\operatorname{ad}_\cdot^*(e_m)e_j^*]-\operatorname{ad}_\cdot^*(\operatorname{ad}_\circ^*(e_j^*)e_m)e_m^*+[\operatorname{ad}_\cdot^*(e_m)e_i^*,e_j^*]\\
&-(\operatorname{ad}_\cdot^*)(e_m)[e_i^*,e_j^*]
\end{aligned}$$

$$= \sum_{l=1}^{n}\sum_{k=1}^{n}\{-(f_{ik}^m+f_{ki}^m)(c_{kl}^j+c_{lk}^j)-(c_{mk}^j+c_{km}^j)(f_{ik}^l+f_{ki}^l)-(f_{jk}^m+f_{kj}^m)(c_{kl}^i+c_{lk}^i)$$
$$-(c_{mk}^i+c_{km}^i)(f_{kj}^l+f_{jk}^l)\}e_l^*.$$

Then, with $\mathcal{G}(\mathcal{A}) \bowtie_{-L_\circ^*}^{-1,-L_\cdot^*} \mathcal{G}(\mathcal{A}^*)$ and the above relation gives

$$\sum_{k=1}^{n}(f_{ij}^k+f_{ji}^k)(c_{ml}^k+c_{lm}^k) = \sum_{k=1}^{n} -(f_{ik}^m+f_{ki}^m)(c_{kl}^j+c_{lk}^j) - (c_{mk}^j+c_{km}^j)(f_{ik}^l+f_{ki}^l)$$
$$-(f_{jk}^m+f_{kj}^m)(c_{kl}^i+c_{lk}^i) - (c_{mk}^i+c_{km}^i)(f_{jk}^l+f_{kj}^l),$$

i.e.

$$\sum_{k=1}^{n}(f_{ij}^k+f_{ji}^k)c_{ml}^k+\sum_{k=1}^{n} c_{kl}^j(f_{ik}^m+f_{ki}^m)+c_{kl}^i(f_{jk}^m+f_{kj}^m)+c_{mk}^j(f_{ik}^l+f_{ki}^l)-c_{mk}^i(f_{jk}^l+f_{kj}^l)$$
$$=\sum_{k=1}^{n}-(f_{ij}^k+f_{ji}^k)c_{lm}^k-\sum_{k=1}^{n}c_{lk}^j(f_{ik}^m+f_{ki}^m)-c_{lk}^i(f_{jk}^m+f_{kj}^m)-c_{km}^j(f_{ik}^l+f_{ki}^l)-c_{km}^i(f_{jk}^l+f_{kj}^l).$$

Replacing l, (resp. m), by m,(resp. l), in the right-hand side of the equality leads to

$$\sum_{k=1}^{n}(f_{ij}^k+f_{ji}^k)c_{ml}^k=\sum_{k=1}^{n}-c_{kl}^j(f_{ik}^m+f_{ki}^m)-c_{kl}^i(f_{jk}^m+f_{kj}^m)-c_{mk}^j(f_{ik}^l+f_{ki}^l)-c_{mk}^i(f_{kj}^l+f_{jk}^l),$$

which is identical to the equation (4.7). □

Definition 4.4. Let $\mathcal{A}$ be a vector space. A *quasi-antiassociative bialgebra* structure on $\mathcal{A}$ is a pair of linear maps (α, β) such that $\alpha : \mathcal{A} \to \mathcal{A} \otimes \mathcal{A}$, $\beta : \mathcal{A}^* \to \mathcal{A}^* \otimes \mathcal{A}^*$ satisfying:

1. $\alpha^* : \mathcal{A}^* \otimes \mathcal{A}^* \to \mathcal{A}^*$ is a quasi-antiassociative algebra structure on $\mathcal{A}^*$,
2. $\beta^* : \mathcal{A} \otimes \mathcal{A} \to \mathcal{A}$ is a quasi-antiassociative algebra structure on $\mathcal{A}$,
3. $\alpha : \mathcal{A} \to \mathcal{A} \otimes \mathcal{A}$ is a 1-cocycle of $\mathcal{G}(\mathcal{A})$ associated to $L_\cdot \otimes 1 + 1 \otimes \mathrm{ad}_\cdot$,
4. $\beta : \mathcal{A}^* \to \mathcal{A}^* \otimes \mathcal{A}^*$ is a 1-cocycle of $\mathcal{G}(\mathcal{A}^*)$ associated to $L_\circ \otimes 1 + 1 \otimes \mathrm{ad}_\circ$.

We also denote this quasi-antiassociative bialgebra by $(\mathcal{A}, \mathcal{A}^*, \alpha, \beta)$ or simply $(\mathcal{A}, \mathcal{A}^*)$.

Remark 4.5. A quasi-antiassociative bialgebra can also be called a Mock-pre-Lie bialgebra or a pre-Jacobi-Jordan bialgebra.

Proposition 4.6. *Let $(\mathcal{A}, \mathcal{A}^*, \alpha, \beta)$ be a quasi-antiassociative bialgebra. Then its dual $(\mathcal{A}^*, \mathcal{A}, \beta, \alpha)$ is also a quasi-antiassociative bialgebra.*

Proposition 4.7. *Let $(\mathcal{A}, \cdot)$ be a quasi-antiassociative algebra and $(\mathcal{A}^*, \circ)$ be a quasi-antiassociative algebra structure on its dual space $\mathcal{A}^*$. Then the following conditions are equivalent:*

1. $(\mathcal{A}, \mathcal{A}^*, \mathrm{ad}^*_\cdot, -R^*_\cdot, \mathrm{ad}^*_\circ, -R^*_\circ)$ *is a matched pair of quasi-antiassociative algebras;*
2. $(\mathcal{G}(\mathcal{A}), \mathcal{G}(\mathcal{A}^*), L^*_\cdot, L^*_\circ)$ *is a matched pair of sub-adjacent Mock-Lie algebras;*
3. $(\mathcal{A}, \mathcal{A}^*)$ *is a quasi-antiassociative bialgebra.*

Definition 4.8. Let $(\mathcal{A}, \mathcal{A}^*, \alpha_\mathcal{A}, \beta_\mathcal{A})$ and $(\mathcal{H}, \mathcal{H}^*, \alpha_\mathcal{H}, \beta_\mathcal{H})$ be two quasi-antiassociative bialgebras. A homomorphism of quasi-antiassociative bialgebras $\varphi : \mathcal{A} \to \mathcal{H}$ is a homomorphism of quasi-antiassociative algebras such that $\varphi^* : \mathcal{H}^* \to \mathcal{A}^*$ is also a homomorphism of quasi-antiassociative algebras, that is, φ satisfies

$$(\varphi\otimes\varphi)\alpha_\mathcal{A}(x) = \alpha_\mathcal{H}(\varphi(x)), \ \ (\varphi^*\otimes\varphi^*)\beta_\mathcal{H}(a^*) = \beta_\mathcal{A}(\varphi^*(a^*)), \forall\, x \in \mathcal{A}, a^* \in \mathcal{H}^*.$$

An isomorphism of quasi-antiassociative bialgebras is an invertible homomorphism of quasi-antiassociative bialgebras.

Proposition 4.9. *Two matched pairs of Mock-Lie algebras are isomorphic if and only if their corresponding quasi-antiassociative bialgebras are isomorphic.*

Proof. Let $(\mathcal{G}(\mathcal{A}) \bowtie \mathcal{G}(\mathcal{A}^*), \mathcal{G}(\mathcal{A}), \mathcal{G}(\mathcal{A}^*))$ and $(\mathcal{G}(\mathcal{H}) \bowtie \mathcal{G}(\mathcal{H}^*), \mathcal{G}(\mathcal{H}), \mathcal{G}(\mathcal{H}^*))$ be two matched pairs of Mock-Lie algebras. Let $\{e_1, \cdots, e_n\}$ be a basis of $\mathcal{A}$ and $\{e_1^*, \cdots, e_n^*\}$ be its dual basis. If $\varphi : \mathcal{G}(\mathcal{A}) \bowtie \mathcal{G}(\mathcal{A}^*) \to \mathcal{G}(\mathcal{H}) \bowtie \mathcal{G}(\mathcal{H}^*)$ is an isomorphism of matched pairs of Mock-Lie algebras, then $\varphi|_\mathcal{A} : \mathcal{A} \to \mathcal{H}$ and $\varphi|_{\mathcal{A}^*} : \mathcal{A}^* \to \mathcal{H}^*$ are isomorphisms of quasi-antiassociative algebras. Moreover, $\varphi|_{\mathcal{A}^*} = (\varphi|_\mathcal{A})^{*^{-1}}$. Hence $(\mathcal{A}, \mathcal{A}^*)$ and $(\mathcal{H}, \mathcal{H}^*)$ are isomorphic as quasi-antiassociative bialgebras. On the other hand, let $(\mathcal{A}, \mathcal{A}^*)$ and $(\mathcal{H}, \mathcal{H}^*)$ be two isomorphic quasi-antiassociative bialgebras and $\varphi' : \mathcal{A} \to \mathcal{H}$ be an isomorphism of quasi-antiassociative bialgebras. Set $\varphi : \mathcal{A} \oplus \mathcal{A}^* \to \mathcal{H} \oplus \mathcal{H}^*$ be a linear map given by

$$\varphi(x) = \varphi'(x), \ \ \varphi(a^*) = (\varphi'^*)^{-1}(a^*), \ \ \forall x \in \mathcal{A}, a^* \in \mathcal{A}^*.$$

Then it is easy to know that φ is an isomorphism of the two matched pairs of Mock-Lie algebras $(\mathcal{G}(\mathcal{A}) \bowtie \mathcal{G}(\mathcal{A}^*), \mathcal{G}(\mathcal{A}), \mathcal{G}(\mathcal{A}^*))$ and $(\mathcal{G}(\mathcal{H}) \bowtie \mathcal{G}(\mathcal{H}^*), \mathcal{G}(\mathcal{H}), \mathcal{G}(\mathcal{H}^*))$. □

Acknowledgment. We sincerely thank Professor Wieslaw Dudek for his valuable suggestions and insightful feedback, particularly regarding the terminology in this work. His thoughtful recommendations have contributed to improving the clarity and consistency of our presentation.

References

[1] **C. Bai**, *Left-symmetric bialgebras and an analogue of the classical Yang-Baxter equation.* J. Commun. Contemp. Math. **2** (2010), 221 – 260.

[2] **A. Baklouti, S. Benayadi, A. Makhlouf, and S. Mansour**, *Cohomology and deformations of Jacobi–Jordan algebras.* ArXiv: 2109.12364v1[math.RA] (2021), 23 pp.

[3] **D. Burde**, *Left-symmetric algebras, or pre-Lie algebras in geometry and physics,* Cent. Eur. J. Math. **4** (2006), 323 – 357.

[4] **M. Camacho, I. Kaygorodov, V. Lopatkin and M.A. Salim**, *The variety of dual mock-Lie algebras.* arXiv:1910.01484 [math.RA].

[5] **F. Chapoton and M. Livernet**, *Pre-Lie algebras and the rooted trees operad,* Int. Math. Res. Not. (2001), 395 – 408.

[6] **M. Gerstenhaber**, *The cohomology structure of an associative ring.* Ann. Math. **78** (1963), 267 – 288.

[7] **T. Kepka**, *Structure of triabelian quasigroups*, Comment. Math. Univ. Carolin. **17** (1976), no. 2, 229 – 240.

[8] **H. Kim**, *Complete left-invariant affine structures on nilpotent Lie groups.* J. Differential Geom. **24** (1986), 373 – 394.

[9] **J.-L. Koszul**, *Domaines bornés homogénes et orbites de groupes de transformations affines.* Bull. Soc. Math. France **89** (1961), 515 – 533.

[10] **M. Markl and E. Remm**, *(Non-)Koszulness of operads for n-ary algebras,galgalim and other curiosities.* J. Homotopy Relat. Struct. **10** (2015), 939 – 969.

[11] **Y. Matsushima**, *Affine structures on complex mainfolds.* Osaka J. Math. **5** (1968), 215 – 222.

[12] **A. Medina**, *Flat left-invariant connections adapted to the automorphism structure of a Lie group.* J. Differential Geom. **16** (1981), 445 – 474.

[13] **S. Okubo and N. Kamiya**, *Jordan-Lie super algebra and Jordan-Lie triple system*, J. Algebra **198** (1997), 388.

[14] **E.B. Vinberg**, *Convex homogeneous cones*, Transl. of Moscow Math. Soc. **12** (1963), 340 – 403.

[15] **P. Zusmanovich**, *Special and exceptional Mock-Lie algebras*, Linear Algebra and its Appl. **518** (2017), 79 – 96.

Received November 21, 2024
Revised April 8, 2025

C.E. Haliya
University of Abomey-Calavi,
International Chair in Mathematical Physics and Applications,
ICMPA-UNESCO Chair, 072 BP 50,
Cotonou, Rep. of Benin
E-mail: cyzille19@gmail.com

G.D. Houndedji, B.Kourouma
Department of Mathematics,
Gamal Abdel Nasser University of Conakry,
Republic of Guinea
E-mail: houndedjidamien@gmail.com, kouroumabakr22@gmail.com